



Heat asymptotics for nonminimal Laplace type operators and application to noncommutative tori

B. Iochum, T. Masson*

Centre de Physique Théorique,
Aix Marseille Univ, Université de Toulon, CNRS, CPT, Marseille, France



ARTICLE INFO

Article history:

Received 25 September 2017

Received in revised form 7 February 2018

Accepted 14 February 2018

Available online 7 March 2018

MSC:

58J35

35J47

81T13

46L87

Keywords:

Heat kernel

Nonminimal operator

Asymptotic heat trace

Laplace type operator

Scalar curvature

Noncommutative torus

ABSTRACT

Let P be a Laplace type operator acting on a smooth hermitean vector bundle V of fiber \mathbb{C}^N over a compact Riemannian manifold given locally by $P = -[g^{\mu\nu}u(x)\partial_\mu\partial_\nu + v^\nu(x)\partial_\nu + w(x)]$ where u, v^ν, w are $M_N(\mathbb{C})$ -valued functions with $u(x)$ positive and invertible. For any $a \in \Gamma(\text{End}(V))$, we consider the asymptotics $\text{Tr}(a e^{-tP}) \underset{t \downarrow 0^+}{\sim} \sum_{r=0}^{\infty} a_r(a, P) t^{(r-d)/2}$ where the coefficients $a_r(a, P)$ can be written as an integral of the functions $a_r(a, P)(x) = \text{tr}[a(x)\mathcal{R}_r(x)]$.

The computation of \mathcal{R}_2 is performed opening the opportunity to calculate the modular scalar curvature for noncommutative tori.

© 2018 Elsevier B.V. All rights reserved.

1. Introduction

As in [1], we consider a d -dimensional compact Riemannian manifold (M, g) without boundary, together with a nonminimal Laplace type operator P on a smooth hermitean vector bundle V over M of fiber \mathbb{C}^N written locally as

$$P := -[g^{\mu\nu}u(x)\partial_\mu\partial_\nu + v^\nu(x)\partial_\nu + w(x)]. \quad (1.1)$$

Here $u(x) \in M_N(\mathbb{C})$ is a positive and invertible matrix valued function and v^ν, w are $M_N(\mathbb{C})$ matrices valued functions. The operator is expressed in a local trivialization of V over an open subset of M which is also a chart on M with coordinates (x^μ) . This trivialization is such that the adjoint for the hermitean metric corresponds to the adjoint of matrices and the trace on endomorphisms on V becomes the usual trace tr on matrices.

For any $a \in \Gamma(\text{End}(V))$, we consider the asymptotics of the heat-trace

$$\text{Tr}(a e^{-tP}) \underset{t \downarrow 0^+}{\sim} \sum_{r=0}^{\infty} a_r(a, P) t^{(r-d)/2} \quad (1.2)$$

* Corresponding author.

E-mail addresses: bruno.iochum@cpt.univ-mrs.fr (B. Iochum), thierry.masson@cpt.univ-mrs.fr (T. Masson).

where Tr is the operator trace. Each coefficient $a_r(a, P)$ can be written as

$$a_r(a, P) = \int_M a_r(a, P)(x) \, \text{dvol}_g(x) \quad (1.3)$$

where $\text{dvol}_g(x) := |g|^{1/2} dx$ with $|g| := \det(g_{\mu\nu})$. The functions $a_r(a, P)(x)$ can be evaluated (various techniques exist for that) and give expressions of the form

$$a_r(a, P)(x) = \text{tr}[a(x)\mathcal{R}_r(x)],$$

where tr is the trace on matrices and \mathcal{R}_r is a (local) section of $\text{End}(V)$. The local section \mathcal{R}_r of $\text{End}(V)$ is uniquely defined by

$$a_r(a, P) = \varphi(a\mathcal{R}_r), \quad (1.4)$$

where

$$\varphi(a) := \int_M \text{tr}[a(x)] \, \text{dvol}_g(x) \quad (1.5)$$

is the natural combined trace on the algebra of sections of $\text{End}(V)$ associated to (M, g) (the integral) and V (the matrix trace). The choice of this trace is not unique, and changing φ changes \mathcal{R}_r . For instance, since M is compact, one can normalize the integral so that the total volume of M is 1, and also the matrix trace such that the trace of the identity matrix is 1. In that case, denoted by $\mathbb{1}$ the identity operator in $\Gamma(\text{End}(V))$, the new combined trace φ_0 satisfies $\varphi_0(\mathbb{1}) = 1$. In Section 5 φ_0 plays an important role since it corresponds to the unique normalized trace on the noncommutative torus algebra. Another possibility for the choice of φ is to use a Riemannian metric on M which is not the tensor g in P , see Remark 2.6.

The aim of this paper is to present a way to compute \mathcal{R}_r by adapting the techniques developed in [1]. These techniques were strongly motivated by a need in physics for explicit computations of $a_r(\mathbb{1}, P)$, see for instance [2,3] and the reference in [1] for the existing results on the mathematical side. The idea behind the computation of \mathcal{R}_2 is to extract the real matrix content of the coefficient a_2 which is related to the scalar curvature of the manifold M .

In Section 2, two formulas are provided for $\mathcal{R}_2(x)$, both in local coordinates (Theorem 2.3) and in a covariant way (Theorem 2.4) in arbitrary dimension and detailed in low dimensions. In Section 3, some direct applications are also provided, for instance to a conformal like transformed Laplacian. Section 4 is devoted to the details of the computations (see also the ancillary Mathematica [4] notebook file [5]).

In Section 5, another application is given in noncommutative geometry. Namely, we compute the conformally perturbed scalar curvature of rational noncommutative tori (NCT). Since at rational values $\theta = p/q$ of the deformation parameter, the algebras of the NCT are isomorphic to the continuous sections of a bundle over the ordinary tori with fiber in $M_q(\mathbb{C})$, they fit perfectly with our previous framework. The irrational case has been widely studied in [6–18]. The results presented in these papers can be written without explicit reference to the parameter θ . In the rational case, our results confirm this property. Moreover, our method gives an alternative which avoids the theory of pseudodifferential calculus on the noncommutative tori introduced by Connes [19] and detailed in [6,20]. In Appendix B, in order to compare to the results in [8, Theorem 5.2] and [11, Theorem 5.4], we perform the change of variables from u to $\ln(u)$ and the change of operators from the left multiplication by u to the conjugation by u , formalized as a substitution lemma (Lemma B.1).

2. The method and the results

In [1], the computation was done for the special case $a = \mathbb{1}$, for which a lot of simplifications can be used under the trace. We now show that the method described there can be adapted almost without any change to compute the quantities \mathcal{R}_r . Moreover, we present the method in a way that reduces the number of steps in the computations, using from the beginning covariant derivatives on the vector bundle V .

2.1. Notations and preliminary results

In order to start with the covariant form of P (see [1, Section A.4]), let us introduce the following notations. We consider a covariant derivative $\nabla_\mu := \partial_\mu + \eta(A_\mu)$, where η is the representation of the Lie algebra of the gauge group of V on any associated vector bundles (mainly V and $\text{End}(V)$ in the following). Let

$$\alpha_\mu := g_{\rho\sigma}(\partial_\mu g^{\rho\sigma}), \quad \alpha^\mu := g^{\mu\nu}\alpha_\nu = g^{\mu\nu}g_{\rho\sigma}(\partial_\nu g^{\rho\sigma}), \quad \beta_\mu := g_{\mu\sigma}(\partial_\rho g^{\rho\sigma}), \quad \beta^\mu := g^{\mu\nu}\beta_\nu = \partial_\nu g^{\mu\nu}.$$

The covariant form of P associated to ∇ (see [1, eq. (A.11)]) is given by

$$\begin{aligned} P &= -(|g|^{-1/2}\nabla_\mu |g|^{1/2}g^{\mu\nu}u\nabla_\nu + p^\mu\nabla_\mu + q) \\ &= -g^{\mu\nu}u\nabla_\mu\nabla_\nu - (p^\nu + g^{\mu\nu}(\nabla_\mu u) - [\frac{1}{2}\alpha^\nu - \beta^\nu]u)\nabla_\nu - q, \end{aligned} \quad (2.1)$$

where the last equality is obtained using $g^{\mu\nu}\frac{1}{2}\partial_\mu \ln|g| + \partial_\mu g^{\mu\nu} = -[\frac{1}{2}\alpha^\nu - \beta^\nu]$. Here, u is as before, and p^μ , q are as v^μ , w from (1.1), except that they transform homogeneously in a change of trivialization of V . All these (local) functions are $M_N(\mathbb{C})$ -valued (as local sections of $\text{End}(V)$), so that η is the adjoint representation:

$$\nabla_\mu u = \partial_\mu u + [A_\mu, u], \quad \nabla_\mu p^\nu = \partial_\mu p^\nu + [A_\mu, p^\nu], \quad \nabla_\mu q = \partial_\mu q + [A_\mu, q].$$

Let us introduce the total covariant derivative $\widehat{\nabla}_\mu$, which combines ∇_μ with the Levi-Civita covariant derivative induced by the metric g . It satisfies

$$\begin{aligned} \widehat{\nabla}_\mu a^v &= \nabla_\mu a^v + \Gamma_{\mu\rho}^v a^\rho = \partial_\mu a^v + [A_\mu, a^v] + \Gamma_{\mu\rho}^v a^\rho, & \widehat{\nabla}_\mu g^{\alpha\beta} &= 0, \\ \widehat{\nabla}_\mu b_\nu &= \nabla_\mu b_\nu - \Gamma_{\mu\nu}^\rho b_\rho = \partial_\mu b_\nu + [A_\mu, b_\nu] - \Gamma_{\mu\nu}^\rho b_\rho, & \widehat{\nabla}_\mu g_{\alpha\beta} &= 0, \end{aligned}$$

for any $\text{End}(V)$ -valued tensors a^v and b_ν , where $\Gamma_{\mu\rho}^v$ are the Christoffel symbols of g . Let us store the following relations:

$$\widehat{\nabla}_\mu u = \nabla_\mu u, \tag{2.2}$$

$$g^{\mu\nu} \widehat{\nabla}_\mu \widehat{\nabla}_\nu u = g^{\mu\nu} (\nabla_\mu \nabla_\nu u - \Gamma_{\mu\nu}^\rho \nabla_\rho u) = g^{\mu\nu} \nabla_\mu \nabla_\nu u - [\frac{1}{2}\alpha^\mu - \beta^\mu] \nabla_\mu u, \tag{2.3}$$

$$\widehat{\nabla}_\mu p^\mu = \nabla_\mu p^\mu - \frac{1}{2} g_{\alpha\beta} (\partial_\mu g^{\alpha\beta}) p^\mu = \nabla_\mu p^\mu - \frac{1}{2} \alpha_\mu p^\mu.$$

Using $\frac{1}{2}\alpha^\rho - \beta^\rho = g^{\mu\nu} \Gamma_{\mu\nu}^\rho$, one then has

$$P = -(g^{\mu\nu} \widehat{\nabla}_\mu u \widehat{\nabla}_\nu + p^v \widehat{\nabla}_v + q) = -g^{\mu\nu} u \widehat{\nabla}_\mu \widehat{\nabla}_\nu - [p^v + g^{\mu\nu} (\widehat{\nabla}_\mu u)] \widehat{\nabla}_\nu - q. \tag{2.4}$$

Notice that in these expressions the total covariant derivative $\widehat{\nabla}_\nu$ (which is the first to act) will never apply to a tensor valued section of V , so that it could be reduced to the covariant derivative ∇_ν .

The writing of P in terms of a covariant derivative ∇ is of course not unique:

Proposition 2.1. *Let $\nabla'_\mu = \nabla_\mu + \eta(\phi_\mu)$ be another covariant derivative on V . Then*

$$P = -(g^{\mu\nu} \widehat{\nabla}'_\mu u \widehat{\nabla}'_\nu + p'^v \widehat{\nabla}'_v + q'), \tag{2.5}$$

with

$$p'^v = p^v - g^{\mu\nu} (u \phi_\mu + \phi_\mu u), \quad q' = q - g^{\mu\nu} (\widehat{\nabla}_\mu u \phi_\nu) + g^{\mu\nu} u \phi_\mu \phi_\nu - p^\mu \phi_\mu. \tag{2.6}$$

In this proposition, ϕ_μ is as p^μ : it transforms homogeneously in a change of trivializations of V .

Proof. This is a direct computation using relations like $\widehat{\nabla}'_\mu u = \widehat{\nabla}_\mu u + [\phi_\mu, u]$ and $\widehat{\nabla}'_\mu \phi_\nu = \widehat{\nabla}_\mu \phi_\nu + [\phi_\mu, \phi_\nu]$ in (2.5) and comparing with (2.4). ■

Corollary 2.2. *There is a unique covariant derivative ∇ such that $p^\mu = 0$. This implies that we can always write P in the reduced form*

$$P = -(g^{\mu\nu} \widehat{\nabla}_\mu u \widehat{\nabla}_\nu + q). \tag{2.7}$$

Proof. The first part of (2.6) can be solved in ϕ_μ for the condition $p'^v = 0$. Indeed, using results in [21], the positivity and invertibility of u implies that for any ν , the equation $u(g^{\mu\nu} \phi_\mu) + (g^{\mu\nu} \phi_\mu)u = p^v$ has a unique solution given by

$$g^{\mu\nu} \phi_\mu = \frac{1}{2} \int_{-\infty}^{+\infty} \frac{u^{it-1/2} p^v u^{-it-1/2}}{\cosh(\pi t)} dt.$$

So, given any covariant derivative to which p^v is associated as in (2.1), we can shift this covariant derivative with the above solution ϕ_μ to impose $p'^v = 0$. ■

This result extends the one in [22, Section 1.2.1], which is a key ingredient of the method used there. In the following, we could have started with P written as in (2.7). But, on one hand, we will see that this is not necessary to get \mathcal{R}_r in terms of u, p^μ, q (at least for $r = 2$). On the other hand, we will see in Section 5 that the covariant derivative which is naturally given by the geometric framework of the rational noncommutative torus does not imply $p^\mu = 0$, and we will then apply directly the most general result. Obviously, it could be possible to first establish our result for the reduced expression (2.7) and then to go to the general result using Proposition 2.1. But this would complicate unnecessarily the presentation of the method and our results.

2.2. The method

The method described in [1] starts with P written as $P = -g^{\mu\nu} u \partial_\mu \partial_\nu - v^\mu \partial_\mu - w$ and leads to $-P(e^{ix\xi} f) = -e^{ix\xi} [H + K + P]f$ where $H = g^{\mu\nu} u \xi_\mu \xi_\nu$ and $K = -i \xi_\mu (v^\mu + 2g^{\mu\nu} u \partial_\nu)$.

This can be generalized for a covariant writing of P . Using (2.1), one gets

$$\begin{aligned} -P(e^{ix\xi} f) &= e^{ix\xi} \left[-g^{\mu\nu} u \xi_\mu \xi_\nu + i \xi_\mu (p^\mu + g^{\mu\nu} (\nabla_\nu u) - [\frac{1}{2}\alpha^\nu - \beta^\nu] u) + 2g^{\mu\nu} u \nabla_\nu \right. \\ &\quad \left. + g^{\mu\nu} u \nabla_\mu \nabla_\nu + (p^v + g^{\mu\nu} (\nabla_\mu u) - [\frac{1}{2}\alpha^\nu - \beta^\nu] u) \nabla_\nu + q \right] f \\ &= -e^{ix\xi} [H + K + P]f, \end{aligned} \tag{2.8}$$

with

$$H := g^{\mu\nu} u_\xi \xi_\mu \xi_\nu, \quad K := -i \xi_\mu (p^\mu + g^{\mu\nu} (\nabla_\nu u) - [\frac{1}{2} \alpha^\mu - \beta^\mu] u + 2g^{\mu\nu} u \nabla_\nu). \tag{2.9}$$

These relations look like the expressions of H and K given above (see [1, eq. (1.6), (1.7)]) with the replacements

$$\partial_\mu \mapsto \nabla_\mu, \quad v^\mu \mapsto p^\mu + g^{\mu\nu} (\nabla_\nu u) - [\frac{1}{2} \alpha^\mu - \beta^\mu] u, \quad w \mapsto q. \tag{2.10}$$

As in [1], we have $\text{Tr}[ae^{-tP}] = \int dx \text{tr}[a(x)K(t, x, x)]$ with

$$K(t, x, x) = \frac{1}{(2\pi)^d} \int d\xi e^{-ix \cdot \xi} (e^{-tP} e^{ix \cdot \xi}) = \frac{1}{(2\pi)^d} \int d\xi e^{-t(H+K+P)} \mathbb{1} = \frac{1}{t^{d/2}} \frac{1}{(2\pi)^d} \int d\xi e^{-H - \sqrt{t}K - tP} \mathbb{1}.$$

Here $\mathbb{1}$ is the constant 1-valued function. Notice that $K(t, x, x)$ is a density, and that $|g|^{-1/2}K(t, x, x)$ is a true function on M . Using the Lebesgue measure dx instead of $d\text{vol}_g(x)$ is convenient to establish the previous relation which uses Fourier transforms (this point has not been emphasized in [1]).

The asymptotics expansion is obtained by the Volterra series

$$e^{A+B} = e^A + \sum_{k=1}^{\infty} \int_{\Delta_k} ds e^{(1-s_1)A} B e^{(s_1-s_2)A} \dots e^{(s_{k-1}-s_k)A} B e^{s_k A}$$

where

$$\Delta_k := \{s = (s_1, \dots, s_k) \in \mathbb{R}_+^k \mid 0 \leq s_k \leq s_{k-1} \leq \dots \leq s_2 \leq s_1 \leq 1\} \text{ and } \Delta_0 := \emptyset \text{ by convention.}$$

For $A = -H$ and $B = -\sqrt{t}K - tP$, one gets

$$e^{-H - \sqrt{t}K - tP} \mathbb{1} = e^{-H} + \sum_{k=1}^{\infty} (-1)^k f_k [(\sqrt{t}K + tP) \otimes \dots \otimes (\sqrt{t}K + tP)] \tag{2.11}$$

with

$$f_k(\xi)[B_1 \otimes \dots \otimes B_k] := \int_{\Delta_k} ds e^{(s_1-1)H(\xi)} B_1 e^{(s_2-s_1)H(\xi)} B_2 \dots B_k e^{-s_k H(\xi)}, \tag{2.12}$$

$$f_0(\xi)[z] := z e^{-H(\xi)},$$

where B_i are matrix-valued differential operators in ∇_μ depending on x and (linearly in) ξ , and $z \in \mathbb{C}$. Collecting the powers of $t^{1/2}$, one gets

$$\text{Tr}(a e^{-tP}) \underset{t \downarrow 0}{\simeq} t^{-d/2} \sum_{r=0}^{\infty} a_r(a, P) t^{r/2}.$$

Each $a_r(a, P)$ contains an integration along ξ , which kills all the terms in odd power in \sqrt{t} since K is linear in ξ while H is quadratic in ξ : $a_{2n+1}(a, P) = 0$ for any $n \in \mathbb{N}$. For instance, the first two non-zero local coefficients are¹

$$a_0(a, P)(x) = \frac{|g|^{-1/2}}{(2\pi)^d} \text{tr}[a(x) \int d\xi e^{-H(x, \xi)}],$$

$$a_2(a, P)(x) = \frac{|g|^{-1/2}}{(2\pi)^d} \text{tr}[a(x) \int d\xi \int_{\Delta_2} ds e^{(s_1-1)H} K e^{(s_2-s_1)H} K e^{-s_2 H}$$

$$- \frac{1}{(2\pi)^d} \text{tr}[a(x) \int d\xi \int_{\Delta_1} ds e^{(s_1-1)H} P e^{-s_1 H}]$$

(remark the coefficient $|g|^{-1/2}$ added here to be compatible with (1.3)).

The strategy to compute these coefficient is twofold. First, we get rid of the ∇_μ 's in the arguments B_i . This is done using [1, Lemma 2.1], which can be applied here since ∇_μ is a derivation: by iteration of the relation

$$f_k(\xi)[B_1 \otimes \dots \otimes B_i \nabla_\mu \otimes \dots \otimes B_k] = \sum_{j=i+1}^k f_k(\xi)[B_1 \otimes \dots \otimes (\nabla_\mu B_j) \otimes \dots \otimes B_k]$$

$$- \sum_{j=i}^k f_{k+1}(\xi)[B_1 \otimes \dots \otimes B_j \otimes (\nabla_\mu H) \otimes B_{j+1} \otimes \dots \otimes B_k], \tag{2.13}$$

¹ Notice the change with convention in [1]: a_r here corresponds to a_r in [1].

we transform each original term into a sum of operators acting on arguments of the form $B_1 \otimes \cdots \otimes B_k = \mathbb{B}_k^{\mu_1 \dots \mu_\ell} \xi_{\mu_1} \dots \xi_{\mu_\ell}$ (for different values of k) where now all the B_i are matrix-valued functions (of x and ξ), or, equivalently, the $\mathbb{B}_k^{\mu_1 \dots \mu_\ell}$ are $M_N(\mathbb{C})^{\otimes k}$ -valued functions (of x only).

The second step of the strategy is to compute the operators applied to the arguments $\mathbb{B}_k^{\mu_1 \dots \mu_\ell}$. They all look like

$$\frac{1}{(2\pi)^d} \int d\xi \xi_{\mu_1} \dots \xi_{\mu_\ell} f_k(\xi) [\mathbb{B}_k^{\mu_1 \dots \mu_\ell}] \in M_N(\mathbb{C}),$$

where the $f_k(\xi)$ are defined by (2.12) and depend only on u through H . As shown in [1], these operators are related to operators $T_{k,p}(x) : M_N(\mathbb{C})^{\otimes k+1} \rightarrow M_N(\mathbb{C})^{\otimes k+1}$ defined by

$$\begin{aligned} T_{k,p}(x) &:= \frac{1}{(2\pi)^d} \int_{\Delta_k} ds \int d\xi \xi_{\mu_1} \dots \xi_{\mu_{2p}} e^{-\|\xi\|^2 C_k(s,u(x))}, \\ T_{0,0}(x) &:= \frac{1}{(2\pi)^d} \int d\xi e^{-\|\xi\|^2 u(x)} \in M_N(\mathbb{C}), \end{aligned} \tag{2.14}$$

where $\|\xi\|^2 := g^{\mu\nu} \xi_\mu \xi_\nu$ and the $C_k(s, A) : M_N(\mathbb{C})^{\otimes k+1} \rightarrow M_N(\mathbb{C})^{\otimes k+1}$ are the operators

$$\begin{aligned} C_k(s, A)[B_0 \otimes B_1 \otimes \cdots \otimes B_k] &= (1 - s_1) B_0 A \otimes B_1 \otimes \cdots \otimes B_k + (s_1 - s_2) B_0 \otimes B_1 A \otimes \cdots \otimes B_k \\ &\quad + \cdots + s_k B_0 \otimes B_1 \otimes \cdots \otimes B_k A. \end{aligned}$$

Denote by $\mathbf{m} : M_N(\mathbb{C})^{\otimes k+1} \rightarrow M_N(\mathbb{C})$, $B_0 \otimes B_1 \otimes \cdots \otimes B_k \mapsto B_0 B_1 \cdots B_k$ the matrix multiplication, then

$$\frac{1}{(2\pi)^d} \int d\xi \xi_{\mu_1} \dots \xi_{\mu_\ell} f_k(\xi) [B_1 \otimes \cdots \otimes B_k] = \mathbf{m} \circ T_{k,p}(x) [B_0 \otimes B_1 \otimes \cdots \otimes B_k],$$

so that each function $a_r(a, P)(x)$ is expressed formally as a sum

$$a_r(a, P)(x) = |g|^{-1/2} \sum \text{tr}[\mathbf{m} \circ T_{k,p}(x)[a(x) \otimes B_1(x) \otimes \cdots \otimes B_k(x)]]. \tag{2.15}$$

This sum comes from the collection of the original terms in K and P producing the power $t^{r/2}$ and the application of [1, Lemma 2.1] i.e. (2.13). This sum relates the r on the LHS to the possible couples (k, p) on the RHS. The B_i are matrix-valued functions (of x) expressed in terms of the original constituents of H, K , and P and their covariant derivatives.

Let us mention here how the procedure introduced in [1] is adapted to the situation where we have the left factor $a(x)$: in [1], the relation between the $T_{k,p}(x)$ and the $f_k(\xi)$ used a trick which consist to add a $B_0 = \mathbb{1}$ argument in front of $B_1 \otimes \cdots \otimes B_k$ (the purpose of the κ map defined in [1]). Here, $\mathbb{1}$ is simply replaced by $a(x)$. But, since

$$\mathbf{m} \circ T_{k,p}(x)[B_0 \otimes B_1(x) \otimes \cdots \otimes B_k(x)] = B_0 \mathbf{m} \circ T_{k,p}(x)[\mathbb{1} \otimes B_1(x) \otimes \cdots \otimes B_k(x)],$$

it is now easy to propose an expression for the factor \mathcal{R}_r as a sum

$$\mathcal{R}_r = |g|^{-1/2} \sum \mathbf{m} \circ T_{k,p}(x)[\mathbb{1} \otimes B_1(x) \otimes \cdots \otimes B_k(x)]. \tag{2.16}$$

One of the main results of [1] is to express the operators $T_{k,p}$ in terms of universal functions through a functional calculus relation involving the spectrum of u (these relations take place at any fixed value of $x \in M$, that we omit from now on). For $r_i > 0$, $\alpha \in \mathbb{R}$, and $k \in \mathbb{N}$, let

$$\begin{aligned} I_{\alpha,k}(r_0, r_1, \dots, r_k) &:= \int_{\Delta_k} ds [(1 - s_1)r_0 + (s_1 - s_2)r_1 + \cdots + s_k r_k]^{-\alpha} \\ &= \int_{\Delta_k} ds [r_0 + s_1(r_1 - r_0) + \cdots + s_k(r_k - r_{k-1})]^{-\alpha}, \end{aligned}$$

so that

$$I_{\alpha,k}(r_0, \dots, r_0) = \frac{1}{k!} r_0^{-\alpha}.$$

In these functions, the arguments $r_i > 0$ are in the spectrum of the positive matrix u .

Denote by $R_i(A) : M_N(\mathbb{C})^{\otimes k+1} \rightarrow M_N(\mathbb{C})^{\otimes k+1}$ the right multiplication on the i th factor

$$R_i(A)[B_0 \otimes B_1 \otimes \cdots \otimes B_k] := B_0 \otimes B_1 \otimes \cdots \otimes B_i A \otimes \cdots \otimes B_k,$$

then

$$T_{k,p} = g_d G(g)_{\mu_1 \dots \mu_{2p}} I_{d/2+p,k}(R_0(u), R_1(u), \dots, R_k(u)),$$

with

$$\begin{aligned}
 g_d &:= \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} d\xi e^{-|\xi|^2} |g(x)|^2 = \frac{|g|^{1/2}}{2^d \pi^{d/2}}, \\
 G(g)_{\mu_1 \dots \mu_{2p}} &:= \frac{1}{(2\pi)^d g_d} \int d\xi \xi_{\mu_1} \dots \xi_{\mu_{2p}} e^{-g^{\alpha\beta} \xi_\alpha \xi_\beta} \\
 &= \frac{1}{2^{2p} p!} \left(\sum_{\rho \in S_{2p}} g^{\mu_\rho(1)\mu_\rho(2)} \dots g^{\mu_\rho(2p-1)\mu_\rho(2p)} \right) = \frac{(2p)!}{2^{2p} p!} g^{(\mu_1 \mu_2 \dots \mu_{2p-1} \mu_{2p})},
 \end{aligned}
 \tag{2.17}$$

where S_{2p} is the symmetric group of permutations on $2p$ elements and the parenthesis in the index of g is the complete symmetrization over all indices. Notice that the factor $|g|^{1/2}$ in g_d simplifies with the factor $|g|^{-1/2}$ in (2.16).

The universal functions $I_{\alpha,k}$ have been studied in [1, Section 3]. They satisfy a recursive formula valid for $1 \neq \alpha \in \mathbb{R}$ and $k \in \mathbb{N}^*$:

$$I_{\alpha,k}(r_0, \dots, r_k) = \frac{1}{(\alpha-1)} (r_{k-1} - r_k)^{-1} [U_{\alpha-1,k-1}(r_0, \dots, r_{k-2}, r_k) - I_{\alpha-1,k-1}(r_0, \dots, r_{k-1})].
 \tag{2.18}$$

It is possible to give some expressions for the $I_{\alpha,k}$ for any (α, k) . They depend on the parity of d . For d even, the main results are that $I_{n,k}$ are Laurent polynomials for $\mathbb{N} \ni n = (d-r)/2 + k \geq k+1$ ($d \geq r+2$) and $k \in \mathbb{N}^*$, while they exhibit a more complicated expression in terms of ln functions for $\mathbb{N} \ni n = (d-r)/2 + k \leq k$ ($d \leq r$). For d odd, the $I_{n,k}$ can be expressed in terms of square roots of the r_i , but without an *a priori* general expression.

The recursive formula (2.18) can be used to write any $I_{\alpha,k}$ appearing in the computation of the operators $T_{k,p}$ in terms of $I_{\alpha-k+1,1}$. The case $\alpha = 1$ appears in dimension $d = 2$: the fundamental spectral function is $I_{1,1}$, and a direct computation shows that

$$I_{1,1}(r_0, r_1) = \frac{\ln r_0 - \ln r_1}{r_0 - r_1}.$$

Using $\frac{x}{e^x - 1} = \sum_{n=0}^{\infty} \frac{B_n}{n!} x^n$, where B_n are the Bernoulli numbers, one gets, with $x = \ln r_0 - \ln r_1$,

$$r_1 I_{1,1}(r_0, r_1) = \sum_{n=0}^{\infty} \frac{B_n}{n!} [\ln r_0 - \ln r_1]^n.$$

A relation between the Bernoulli numbers and $a_2(a, P)$ has already been noticed in the computation of the modular curvature for the noncommutative two torus in [7] (see Section 5).

2.3. The results for $a_2(a, P)$

In the following, we restrict ourselves to the computation of $a_2(a, P)$. This section gives the main results of the paper. The computations are detailed in Section 4.

Let us introduce the following notation. For any $x \in M$, denote by $r_i = r_i(x) > 0$ an element in the (discrete) spectrum $\text{sp}(u)$ of $u = u(x)$ and by $E_{r_i} = E_{r_i}(x)$ the associated projection of u . This implies that

$$u = \sum_{r_0 \in \text{sp}(u)} r_0 E_{r_0} = r_0 E_{r_0}$$

where in the last expression we omit the summation over r_0 , as will be the case in many expressions given in the following. Notice that $\mathbf{1} = \sum_{r_0 \in \text{sp}(u)} E_{r_0}$ and $E_{r_0} E_{r_1} = \delta_{r_0, r_1} E_{r_0}$.

Theorem 2.3. For P given by (1.1), $a_2(a, P)(x) = \text{tr}[a(x)\mathcal{R}_2(x)]$ with

$$\begin{aligned}
 \mathcal{R}_2 &= \frac{1}{2^d \pi^{d/2}} \left[c r_0^{-d/2+1} E_{r_0} + F_{\partial u}^\mu(r_0, r_1) E_{r_0} (\partial_\mu u) E_{r_1} \right. \\
 &\quad + g^{\mu\nu} F_{\partial \partial u}(r_0, r_1) E_{r_0} (\partial_\mu \partial_\nu u) E_{r_1} + g^{\mu\nu} F_{\partial u, \partial u}(r_0, r_1, r_2) E_{r_0} (\partial_\mu u) E_{r_1} (\partial_\nu u) E_{r_2} \\
 &\quad + F_w(r_0, r_1) E_{r_0} w E_{r_1} + F_{v, \mu}(r_0, r_1) E_{r_0} v^\mu E_{r_1} \\
 &\quad + F_{v, \partial u}(r_0, r_1, r_2) E_{r_0} v^\mu E_{r_1} (\partial_\mu u) E_{r_2} + F_{\partial u, v}(r_0, r_1, r_2) E_{r_0} (\partial_\mu u) E_{r_1} v^\mu E_{r_2} \\
 &\quad \left. + g_{\mu\nu} F_{v, v}(r_0, r_1, r_2) E_{r_0} v^\mu E_{r_1} v^\nu E_{r_2} + F_{\partial v}(r_0, r_1) E_{r_0} (\partial_\mu v^\mu) E_{r_1} \right],
 \end{aligned}
 \tag{2.19}$$

where the sums over the r_0, r_1, r_2 in the spectrum of u are omitted, the spectral functions F are given below, and

$$\begin{aligned}
 c &:= \frac{1}{3} (\partial_\mu \partial_\nu g^{\mu\nu}) - \frac{1}{12} g^{\mu\nu} g_{\rho\sigma} (\partial_\mu \partial_\nu g^{\rho\sigma}) + \frac{1}{48} g^{\mu\nu} g_{\rho\sigma} g_{\alpha\beta} (\partial_\mu g^{\rho\sigma}) (\partial_\nu g^{\alpha\beta}) \\
 &\quad + \frac{1}{24} g^{\mu\nu} g_{\rho\sigma} g_{\alpha\beta} (\partial_\mu g^{\rho\alpha}) (\partial_\nu g^{\sigma\beta}) - \frac{1}{12} g_{\rho\sigma} (\partial_\mu g^{\mu\nu}) (\partial_\nu g^{\rho\sigma}) \\
 &\quad + \frac{1}{12} g_{\rho\sigma} (\partial_\mu g^{\nu\rho}) (\partial_\nu g^{\mu\sigma}) - \frac{1}{4} g_{\rho\sigma} (\partial_\mu g^{\mu\rho}) (\partial_\nu g^{\nu\sigma}).
 \end{aligned}$$

The coefficient c is given here in an arbitrary coordinate system. Since (2.19) is not given in terms of (Riemannian) covariant quantities, c is not expected to have a good behavior under change of coordinates. In normal coordinates, c reduces to the first two terms and is equal to $-R/6$ where R is the scalar curvature. A covariant approach will be given in Theorem 2.4.

The spectral functions in (2.19) are given in terms of the universal function $I_{d/2,1}$ by

$$\begin{aligned}
 F_w(r_0, r_1) &= I_{d/2,1}(r_0, r_1), \\
 F_{\partial v}(r_0, r_1) &= 2r_0 \frac{I_{d/2,1}(r_0, r_0) - I_{d/2,1}(r_0, r_1)}{d(r_0 - r_1)}, \\
 F_{\partial\partial u}(r_0, r_1) &= -r_0 \frac{4r_0 I_{d/2,1}(r_0, r_0) + ((d-4)r_0 - dr_1) I_{d/2,1}(r_0, r_1)}{d(r_0 - r_1)^2}, \\
 F_{\partial u}^\mu(r_0, r_1) &= [\frac{1}{2}\alpha^\mu - \beta^\mu] r_0 \frac{4r_0 I_{d/2,1}(r_0, r_0) + ((d-4)r_0 - dr_1) I_{d/2,1}(r_0, r_1)}{d(r_0 - r_1)^2}, \\
 F_{v,\mu}(r_0, r_1) &= -\alpha_\mu r_0 \frac{I_{d/2,1}(r_0, r_0) - I_{d/2,1}(r_0, r_1)}{d(r_0 - r_1)} - \frac{1}{2}[\frac{1}{2}\alpha_\mu - \beta_\mu] I_{d/2,1}(r_0, r_1), \\
 F_{v,v}(r_0, r_1, r_2) &= \frac{I_{d/2,1}(r_0, r_1) - I_{d/2,1}(r_0, r_2)}{d(r_1 - r_2)}, \\
 F_{\partial u,v}(r_0, r_1, r_2) &= \frac{2r_0}{d} \left[\frac{I_{d/2,1}(r_0, r_0)}{(r_0 - r_1)(r_0 - r_2)} + \frac{I_{d/2,1}(r_0, r_1)}{(r_1 - r_0)(r_1 - r_2)} + \frac{I_{d/2,1}(r_0, r_2)}{(r_2 - r_0)(r_2 - r_1)} \right], \\
 F_{v,\partial u}(r_0, r_1, r_2) &= -2r_0 \frac{I_{d/2,1}(r_0, r_0)}{d(r_0 - r_2)(r_1 - r_2)} - 2r_1 \frac{I_{d/2,1}(r_0, r_1)}{d(r_1 - r_2)^2} \\
 &\quad - \frac{((d-4)r_0 r_1 - (d-2)r_0 r_2 - (d-2)r_1 r_2 + dr_1^2) I_{d/2,1}(r_0, r_2)}{d(r_0 - r_2)(r_1 - r_2)^2}, \\
 F_{\partial u,\partial u}(r_0, r_1, r_2) &= \frac{4r_0}{d(r_0 - r_1)(r_0 - r_2)^2(r_1 - r_2)^2} \\
 &\quad \times [r_0(r_1 - r_2)(r_0 - 2r_1 + r_2) I_{d/2,1}(r_0, r_0) + r_1(r_0 - r_2)^2 I_{d/2,1}(r_0, r_1) \\
 &\quad + \frac{1}{2}(r_0 - r_1)((d-4)r_0 r_1 - (d-2)r_0 r_2 - dr_1 r_2 + (d+2)r_1^2) I_{d/2,1}(r_0, r_2)].
 \end{aligned}$$

Theorem 2.4. For P given by (2.1), $a_2(a, P)(x) = \text{tr}[a(x)\mathcal{R}_2(x)]$ with

$$\begin{aligned}
 \mathcal{R}_2 &= \frac{1}{2^d \pi^{d/2}} \left[\frac{1}{6} R r_0^{-d/2+1} E_{r_0} + G_q(r_0, r_1) E_{r_0} q E_{r_1} + g^{\mu\nu} G_{\widehat{\nabla}\widehat{\nabla}u}(r_0, r_1) E_{r_0} (\widehat{\nabla}_\mu \widehat{\nabla}_\nu u) E_{r_1} \right. \\
 &\quad + G_{\widehat{\nabla}p}(r_0, r_1) E_{r_0} (\widehat{\nabla}_\mu p^\mu) E_{r_1} + g^{\mu\nu} G_{\widehat{\nabla}u, \widehat{\nabla}u}(r_0, r_1, r_2) E_{r_0} (\widehat{\nabla}_\mu u) E_{r_1} (\widehat{\nabla}_\nu u) E_{r_2} \\
 &\quad + G_{p, \widehat{\nabla}u}(r_0, r_1, r_2) E_{r_0} p^\mu E_{r_1} (\widehat{\nabla}_\mu u) E_{r_2} + G_{\widehat{\nabla}u, p}(r_0, r_1, r_2) E_{r_0} (\widehat{\nabla}_\mu u) E_{r_1} p^\mu E_{r_2} \\
 &\quad \left. + G_{p,p}(r_0, r_1, r_2) E_{r_0} p^\mu E_{r_1} p_\mu E_{r_2} \right] \tag{2.20}
 \end{aligned}$$

where the sums over the r_0, r_1, r_2 in the spectrum of u are omitted, the spectral functions G are given below, and R is the scalar curvature of g .

The spectral functions in (2.20) are given in terms of the spectral functions F by

$$\begin{aligned}
 G_q(r_0, r_1) &:= F_w(r_0, r_1) \\
 &= G_q(r_1, r_0), \\
 G_{\widehat{\nabla}\widehat{\nabla}u}(r_0, r_1) &:= F_{\partial\partial u}(r_0, r_1) + F_{\partial v}(r_0, r_1) \\
 &= G_{\widehat{\nabla}\widehat{\nabla}u}(r_1, r_0), \\
 G_{\widehat{\nabla}p}(r_0, r_1) &:= F_{\partial v}(r_0, r_1), \\
 G_{\widehat{\nabla}u, \widehat{\nabla}u}(r_0, r_1, r_2) &:= F_{\partial u, \partial u}(r_0, r_1, r_2) + F_{v, \partial u}(r_0, r_1, r_2) + F_{\partial u, v}(r_0, r_1, r_2) + F_{v, v}(r_0, r_1, r_2) \\
 &= G_{\widehat{\nabla}u, \widehat{\nabla}u}(r_2, r_1, r_0), \\
 G_{p, \widehat{\nabla}u}(r_0, r_1, r_2) &:= F_{v, \partial u}(r_0, r_1, r_2) + F_{v, v}(r_0, r_1, r_2), \\
 G_{\widehat{\nabla}u, p}(r_0, r_1, r_2) &:= F_{\partial u, v}(r_0, r_1, r_2) + F_{v, v}(r_0, r_1, r_2) \\
 &= -G_{p, \widehat{\nabla}u}(r_2, r_1, r_0), \\
 G_{p,p}(r_0, r_1, r_2) &:= F_{v, v}(r_0, r_1, r_2) \\
 &= G_{p,p}(r_2, r_1, r_0).
 \end{aligned}$$

As shown in [1], the universal spectral functions $I_{\alpha,k}$ are continuous, so that all the spectral functions F and G are also continuous, as can be deduced from their original expressions in terms of functions $I_{\alpha,k}$ given in the list (4.1) and the above relations between the F and the G .

Remark 2.5 (Homogeneity by Dilation). Using (1.2) and (1.4), we get $\mathcal{R}_r(\lambda P) = \lambda^{(r-d)/2} \mathcal{R}_r(P)$ for any $\lambda \in \mathbb{R}_+^*$. The dilation of P by λ is equivalent to the dilations of u, v^μ, w, p^μ, q by λ . Using $I_{\alpha,k}(\lambda r_0, \dots, \lambda r_k) = \lambda^{-\alpha} I_{\alpha,k}(r_0, \dots, r_k)$ and the explicit expressions of the spectral functions, all terms in (2.19) and (2.20) are λ -homogeneous of degree $(2-d)/2$. \square

Remark 2.6. The metric g plays a double role here: it is the metric of the Riemannian manifold (M, g) and it is the non-degenerate tensor which multiply u in P . If one has to consider two operators P_1 and P_2 with tensors g_1 and g_2 on the same manifold M , it may be not natural to take g_1 or g_2 as the Riemannian metric on M . It is possible to consider a metric h on M different to the tensor g associated to P . In that case, we have to replace dvol_g in (1.3) by dvol_h and $K(t, x, x, \cdot)$ is then a density for h , so that the true function is now $|h|^{-1/2} K(t, x, x)$, and $|h|^{-1/2}$ appears in (2.15) and (2.16) in place of $|g|^{-1/2}$. Now, the computation of $T_{k,p}$ makes apparent the coefficient g_d given by (2.17) where the metric g comes from P . Finally, in (2.19) and (2.20), the two determinants do not simplify anymore, and one gets an extra factor $|g|^{1/2} |h|^{-1/2}$ in front of \mathcal{R}_2 , which is now relative to $\varphi_h(a) := \int_M \text{tr}[a(x)] \text{dvol}_h(x)$. \square

A change of connection as in Proposition 2.1 does not change the value of \mathcal{R}_2 . This induces the following relations between the spectral functions G .

Proposition 2.7. *The spectral functions G satisfy the relations:*

$$\begin{aligned} G_{\widehat{\nabla}_p}(r_0, r_1) &= -\frac{r_0 G_q(r_0, r_1) + (r_0 - r_1) G_{\widehat{\nabla}_p \widehat{\nabla}_u}(r_0, r_1)}{r_0 + r_1}, \\ G_{\widehat{\nabla}_{u,p}}(r_0, r_1, r_2) &= -\frac{r_2 G_q(r_0, r_2) + (r_0 + 3r_2) G_{\widehat{\nabla}_p \widehat{\nabla}_u}(r_0, r_2) + (r_0 + r_2)(r_1 - r_2) G_{\widehat{\nabla}_u, \widehat{\nabla}_u}(r_0, r_1, r_2)}{(r_0 + r_2)(r_1 + r_2)}, \\ G_{p, \widehat{\nabla}_u}(r_0, r_1, r_2) &= \frac{r_0 G_q(r_0, r_2) + (3r_0 + r_2) G_{\widehat{\nabla}_p \widehat{\nabla}_u}(r_0, r_2) + (r_0 + r_2)(r_1 - r_0) G_{\widehat{\nabla}_u, \widehat{\nabla}_u}(r_0, r_1, r_2)}{(r_0 + r_2)(r_1 + r_0)}, \\ G_{p,p}(r_0, r_1, r_2) &= -\frac{r_1 G_q(r_0, r_2) - (r_0 - 2r_1 + r_2) G_{\widehat{\nabla}_p \widehat{\nabla}_u}(r_0, r_2) + (r_0 - r_1)(r_1 - r_2) G_{\widehat{\nabla}_u, \widehat{\nabla}_u}(r_0, r_1, r_2)}{(r_0 + r_1)(r_1 + r_2)}. \end{aligned}$$

Proof. Inserting the relations (2.6) into (2.20), all the terms involving ϕ_μ must vanish. This induces the following relations between the G functions:

$$\begin{aligned} r_0 G_q(r_0, r_1) + (r_0 - r_1) G_{\widehat{\nabla}_p \widehat{\nabla}_u}(r_0, r_1) + (r_0 + r_1) G_{\widehat{\nabla}_p}(r_0, r_1) &= 0, \\ G_{\widehat{\nabla}_p}(r_0, r_2) - (r_0 - r_1) G_{\widehat{\nabla}_{u,p}}(r_0, r_1, r_2) - (r_0 + r_1) G_{p,p}(r_0, r_1, r_2) &= 0, \\ G_q(r_0, r_2) + G_{\widehat{\nabla}_p}(r_0, r_2) + (r_1 - r_2) G_{p, \widehat{\nabla}_u}(r_0, r_1, r_2) + (r_1 + r_2) G_{p,p}(r_0, r_1, r_2) &= 0, \\ 2G_{\widehat{\nabla}_p \widehat{\nabla}_u}(r_0, r_2) - G_{\widehat{\nabla}_p}(r_0, r_2) - (r_0 - r_1) G_{\widehat{\nabla}_u, \widehat{\nabla}_u}(r_0, r_1, r_2) - (r_0 + r_1) G_{p, \widehat{\nabla}_u}(r_0, r_1, r_2) &= 0, \\ G_q(r_0, r_2) + 2G_{\widehat{\nabla}_p \widehat{\nabla}_u}(r_0, r_2) + G_{\widehat{\nabla}_p}(r_0, r_2) &+ (r_1 - r_2) G_{\widehat{\nabla}_u, \widehat{\nabla}_u}(r_0, r_1, r_2) + (r_1 + r_2) G_{\widehat{\nabla}_{u,p}}(r_0, r_1, r_2) = 0, \\ r_0 G_q(r_0, r_2) + (r_0 - 2r_1 + r_2) G_{\widehat{\nabla}_p \widehat{\nabla}_u}(r_0, r_2) + (r_0 - r_2) G_{\widehat{\nabla}_p}(r_0, r_2) &+ (r_0 - r_1)(r_1 - r_2) G_{\widehat{\nabla}_u, \widehat{\nabla}_u}(r_0, r_1, r_2) + (r_0 + r_1)(r_1 - r_2) G_{p, \widehat{\nabla}_u}(r_0, r_1, r_2) \\ + (r_0 - r_1)(r_1 + r_2) G_{\widehat{\nabla}_{u,p}}(r_0, r_1, r_2) + (r_0 + r_1)(r_1 + r_2) G_{p,p}(r_0, r_1, r_2) &= 0. \end{aligned}$$

One can check directly that these relations hold true. From them, one can solve $G_{\widehat{\nabla}_p}$, $G_{\widehat{\nabla}_{u,p}}$, $G_{p, \widehat{\nabla}_u}$, and $G_{p,p}$ in terms of G_q , $G_{\widehat{\nabla}_p \widehat{\nabla}_u}$, and $G_{\widehat{\nabla}_u, \widehat{\nabla}_u}$. This gives the relations of the proposition. \blacksquare

These relations show that the four spectral functions G involved in terms with p^μ are deduced from the three spectral functions involving only u and q . This result is not a surprise: from Corollary 2.2 we know that we can start with $p^\mu = 0$, so that \mathcal{R}_2 is written in terms of the three functions G_q , $G_{\widehat{\nabla}_p \widehat{\nabla}_u}$, $G_{\widehat{\nabla}_u, \widehat{\nabla}_u}$ only, and then we can change the connection in order to produce the most general expression for \mathcal{R}_2 . In other words, among the seven spectral functions G , only three are fundamental.

The spectral functions G can be computed explicitly, and their expressions depend on the value of m . For $d = 2$ and $d = 3$, these spectral functions are written in terms of the following functions:

$$\begin{aligned} Q_1(a, b, c) &:= \frac{-3a^3 + a^2b - 6a^2c + 6abc + ac^2 + bc^2}{2(a-b)^2(a-c)^3}, \\ Q_2(a, b, c) &:= \frac{1}{2} \left[\frac{2}{(a-b)(a-c)} + \frac{a+b}{(b-a)^2(b-c)} \ln(b/a) + \frac{a+c}{(c-a)^2(c-b)} \ln(c/a) \right], \\ Q_3(a, b, c) &:= 6\sqrt{abc} + a^{3/2} + 2b^{3/2} + c^{3/2} + \sqrt{ac}(\sqrt{a} + \sqrt{c}) \\ &+ 2(a(\sqrt{b} + \sqrt{c}) + 2b(\sqrt{a} + \sqrt{c}) + c(\sqrt{a} + \sqrt{b})), \\ Q_4(a, b, c) &:= \frac{a+b+c + 2\sqrt{ab} + 2\sqrt{ac} + \sqrt{bc}}{\sqrt{bc}(\sqrt{a} + \sqrt{b})^2(\sqrt{a} + \sqrt{c})^2(\sqrt{b} + \sqrt{c})}. \end{aligned}$$

Corollary 2.8 (Case $d = 2$). In dimension two the spectral functions G can be written in terms of log functions:

$$\begin{aligned} \mathcal{R}_2 = & \frac{1}{4\pi} \left[\frac{1}{6}R + \frac{\ln(r_0/r_1)}{r_0 - r_1} E_{r_0} q E_{r_1} + \frac{1}{r_0 - r_1} \left(1 - r_0 \frac{\ln(r_0/r_1)}{r_0 - r_1}\right) E_{r_0} (\widehat{\nabla}_\mu p^\mu) E_{r_1} \right. \\ & - \frac{1}{(r_0 - r_1)^2} (r_0 + r_1 - 2r_0 r_1 \frac{\ln(r_0/r_1)}{r_0 - r_1}) g^{\mu\nu} E_{r_0} (\widehat{\nabla}_\mu \widehat{\nabla}_\nu u) E_{r_1} \\ & + \left(\frac{(r_0 + r_2)(r_0 - 2r_1 + r_2)}{(r_0 - r_1)(r_0 - r_2)^2(r_1 - r_2)} \right. \\ & \quad \left. - Q_1(r_0, r_1, r_2) \ln(r_0/r_1) - Q_1(r_2, r_1, r_0) \ln(r_2/r_1) \right) g^{\mu\nu} E_{r_0} (\widehat{\nabla}_\mu u) E_{r_1} (\widehat{\nabla}_\nu u) E_{r_2} \\ & + Q_2(r_0, r_1, r_2) E_{r_0} (\widehat{\nabla}_\mu u) E_{r_1} p^\mu E_{r_2} - Q_2(r_2, r_1, r_0) E_{r_0} p^\mu E_{r_1} (\widehat{\nabla}_\mu u) E_{r_2} \\ & \left. + \frac{(r_1 - r_2) \ln(r_0/r_1) + (r_0 - r_1) \ln(r_2/r_1)}{2(r_0 - r_1)(r_0 - r_2)(r_1 - r_2)} E_{r_0} p_\mu E_{r_1} p^\mu E_{r_2} \right]. \end{aligned}$$

When $d = 2m \geq 4$ is even, all the involved functions are Laurent polynomials as a direct consequence of [1, Prop. 3.5]:

$$\begin{aligned} G_q(r_0, r_1) &= \sum_{0 \leq \ell \leq m-2} \frac{1}{m-1} r_0^{\ell+1-m} r_1^{-\ell-1}, \\ G_{\widehat{\nabla}\widehat{\nabla}u}(r_0, r_1) &= - \sum_{0 \leq \ell \leq m-2} \frac{(m-\ell-1)(\ell+1)}{m(m-1)} r_0^{\ell+1-m} r_1^{-\ell-1}, \\ G_{\widehat{\nabla}p}(r_0, r_1) &= - \sum_{0 \leq \ell \leq m-2} \frac{m-\ell-1}{m(m-1)} r_0^{\ell+1-m} r_1^{-\ell-1}, \\ G_{\widehat{\nabla}u, \widehat{\nabla}u}(r_0, r_1, r_2) &= - \sum_{0 \leq \ell \leq k \leq m-2} \frac{(2\ell+1)(2k-2m+3)}{2m(m-1)} r_0^{k+1-m} r_1^{\ell-k-1} r_2^{-\ell-1}, \\ G_{p, \widehat{\nabla}u}(r_0, r_1, r_2) &= - \sum_{0 \leq \ell \leq k \leq m-2} \frac{2\ell+1}{2m(m-1)} r_0^{k+1-m} r_1^{\ell-k-1} r_2^{-\ell-1}, \\ G_{\widehat{\nabla}u, p}(r_0, r_1, r_2) &= - \sum_{0 \leq \ell \leq k \leq m-2} \frac{2k-2m+3}{2m(m-1)} r_0^{k+1-m} r_1^{\ell-k-1} r_2^{-\ell-1}, \\ G_{p, p}(r_0, r_1, r_2) &= - \sum_{0 \leq \ell \leq k \leq m-2} \frac{1}{2m(m-1)} r_0^{k+1-m} r_1^{\ell-k-1} r_2^{-\ell-1}. \end{aligned}$$

This implies the following expressions for \mathcal{R}_2 :

Corollary 2.9 (Case $d = 2m$ Even and $d \geq 4$). Using the expressions of the spectral functions G as Laurent polynomials, one has

$$\begin{aligned} \mathcal{R}_2 = & \frac{1}{2^{2m}\pi^m} \left[\frac{1}{6}R u^{-m+1} + \sum_{0 \leq \ell \leq m-2} \frac{1}{m-1} u^{\ell+1-m} q u^{-\ell-1} - \sum_{0 \leq \ell \leq m-2} \frac{m-\ell-1}{m(m-1)} u^{\ell+1-m} (\widehat{\nabla}_\mu p^\mu) u^{-\ell-1} \right. \\ & - \sum_{0 \leq \ell \leq m-2} \frac{(m-\ell-1)(\ell+1)}{m(m-1)} g^{\mu\nu} u^{\ell+1-m} (\widehat{\nabla}_\mu \widehat{\nabla}_\nu u) u^{-\ell-1} \\ & - \sum_{0 \leq \ell \leq k \leq m-2} \frac{(2\ell+1)(2k-2m+3)}{2m(m-1)} g^{\mu\nu} u^{k+1-m} (\widehat{\nabla}_\mu u) u^{\ell-k-1} (\widehat{\nabla}_\nu u) u^{-\ell-1} \\ & - \sum_{0 \leq \ell \leq k \leq m-2} \frac{2k-2m+3}{2m(m-1)} u^{k+1-m} (\widehat{\nabla}_\mu u) u^{\ell-k-1} p^\mu u^{-\ell-1} \\ & - \sum_{0 \leq \ell \leq k \leq m-2} \frac{2\ell+1}{2m(m-1)} u^{k+1-m} p^\mu u^{\ell-k-1} (\widehat{\nabla}_\mu u) u^{-\ell-1} \\ & \left. - \sum_{0 \leq \ell \leq k \leq m-2} \frac{1}{2m(m-1)} u^{k+1-m} p_\mu u^{\ell-k-1} p^\mu u^{-\ell-1} \right] \\ = & \frac{1}{2^{2m}\pi^m} \left[\frac{1}{6}R u^{-m+1} + \sum_{0 \leq \ell \leq m-2} \frac{1}{m-1} u^{\ell+1-m} q u^{-\ell-1} \right. \\ & - \sum_{0 \leq \ell \leq m-2} \frac{(m-\ell-1)}{m(m-1)} g^{\mu\nu} u^{\ell+1-m} [\widehat{\nabla}_\mu ((\ell+1)\widehat{\nabla}_\nu u + p_\nu)] u^{-\ell-1} \\ & \left. - \sum_{0 \leq \ell \leq k \leq m-2} \frac{1}{2m(m-1)} g^{\mu\nu} u^{k+1-m} [(2k-2m+3)\widehat{\nabla}_\mu u + p_\mu] u^{\ell-k-1} [(2\ell+1)\widehat{\nabla}_\nu u + p_\nu] u^{-\ell-1} \right]. \end{aligned}$$

Corollary 2.10 (Case $d = 4$). In dimension four these expressions simplify further to:

$$\begin{aligned} \mathcal{R}_2 &= \frac{1}{16\pi^2} \left[\frac{1}{6} R u^{-1} + u^{-1} q u^{-1} - \frac{1}{2} u^{-1} (\widehat{\nabla}_\mu p^\mu) u^{-1} - \frac{1}{2} g^{\mu\nu} u^{-1} (\widehat{\nabla}_\mu \widehat{\nabla}_\nu u) u^{-1} \right. \\ &\quad \left. + \frac{1}{4} g^{\mu\nu} u^{-1} (\widehat{\nabla}_\mu u) u^{-1} (\widehat{\nabla}_\nu u) u^{-1} + \frac{1}{4} u^{-1} (\widehat{\nabla}_\mu u) u^{-1} p^\mu u^{-1} - \frac{1}{4} u^{-1} p^\mu u^{-1} (\widehat{\nabla}_\mu u) u^{-1} \right. \\ &\quad \left. - \frac{1}{4} u^{-1} p_\mu u^{-1} p^\mu u^{-1} \right] \\ &= \frac{1}{16\pi^2} \left[\frac{1}{6} R u^{-1} + u^{-1} q u^{-1} - \frac{1}{2} g^{\mu\nu} u^{-1} [\widehat{\nabla}_\mu (\widehat{\nabla}_\nu u + p_\nu)] u^{-1} + \frac{1}{4} g^{\mu\nu} u^{-1} [\widehat{\nabla}_\mu u - p_\mu] u^{-1} [\widehat{\nabla}_\nu u + p_\nu] u^{-1} \right]. \end{aligned}$$

Corollary 2.11 (Case $d = 3$). In dimension three the spectral functions G can be written in terms of square roots of the r_i (see [1, Prop. 3.4]) and this leads to:

$$\begin{aligned} \mathcal{R}_2 &= \frac{1}{8\pi^{3/2}} \left[\frac{1}{6} R + \frac{2}{\sqrt{r_0 r_1} (\sqrt{r_0} + \sqrt{r_1})} E_{r_0} q E_{r_1} - \frac{2}{3} \frac{2\sqrt{r_0} + \sqrt{r_1}}{\sqrt{r_0 r_1} (\sqrt{r_0} + \sqrt{r_1})^2} E_{r_0} (\widehat{\nabla}_\mu p^\mu) E_{r_1} \right. \\ &\quad \left. - \frac{2}{3} \frac{\sqrt{r_0 r_1} + (\sqrt{r_0} + \sqrt{r_1})^2}{\sqrt{r_0 r_1} (\sqrt{r_0} + \sqrt{r_1})^3} g^{\mu\nu} E_{r_0} (\widehat{\nabla}_\mu \widehat{\nabla}_\nu u) E_{r_1} \right. \\ &\quad \left. + \frac{2}{3} \frac{Q_3(r_0, r_1, r_2)}{\sqrt{r_1} (\sqrt{r_0} + \sqrt{r_1})^2 (\sqrt{r_0} + \sqrt{r_2})^3 (\sqrt{r_1} + \sqrt{r_2})^2} g^{\mu\nu} E_{r_0} (\widehat{\nabla}_\mu u) E_{r_1} (\widehat{\nabla}_\nu u) E_{r_2} \right. \\ &\quad \left. + \frac{2}{3} Q_4(r_0, r_1, r_2) E_{r_0} (\widehat{\nabla}_\mu u) E_{r_1} p^\mu E_{r_2} - \frac{2}{3} Q_4(r_2, r_1, r_0) E_{r_0} p_\mu E_{r_1} p^\mu E_{r_2} \right. \\ &\quad \left. - \frac{2}{3} \frac{\sqrt{r_0} + \sqrt{r_1} + \sqrt{r_2}}{\sqrt{r_0 r_1 r_2} (\sqrt{r_0} + \sqrt{r_1}) (\sqrt{r_0} + \sqrt{r_2}) (\sqrt{r_1} + \sqrt{r_2})} E_{r_0} p_\mu E_{r_1} p^\mu E_{r_2} \right]. \end{aligned}$$

To conclude this list of results given at various dimensions, we see that we have explicit expressions of \mathcal{R}_2 for d even, and moreover simple generic expressions for $d = 2m, d \geq 4$, while for d odd it is difficult to propose a generic expression.

3. Some direct applications

3.1. The case $a = \mathbb{1}$

When $a = \mathbb{1}$, for any $k \geq 1$, a cyclic operation can be performed under the trace:

$$\begin{aligned} f(r_0, r_1, \dots, r_k) \operatorname{tr}(E_{r_0} B_1 E_{r_1} B_2 \cdots B_k E_{r_k}) &= f(r_0, r_1, \dots, r_k) \operatorname{tr}(E_{r_k} E_{r_0} B_1 E_{r_1} B_2 \cdots B_k) \\ &= \delta_{r_0, r_k} f(r_0, r_1, \dots, r_k) \operatorname{tr}(E_{r_k} E_{r_0} B_1 E_{r_1} B_2 \cdots B_k) \\ &= f(r_0, r_1, \dots, r_0) \operatorname{tr}(E_{r_0} B_1 E_{r_1} B_2 \cdots E_{r_{k-1}} B_k). \end{aligned}$$

This implies that in all the spectral functions $f(r_0, r_1, \dots, r_k)$ one can put $r_k = r_0$ (remember that all the spectral functions are continuous, so that $r_k \rightarrow r_0$ is well-defined).

In [1, Theorem 4.3], $a_2(\mathbb{1}, P)$ has been computed for $d = 2m$ even, $m \geq 1$. Let us first rewrite this result in terms of spectral functions:

$$\begin{aligned} &c r_0^{-m+1} E_0 + r_0^{-m} E_0 w + \frac{m-2}{6} [\frac{1}{2} \alpha^\mu - \beta^\mu] r_0^{-m} E_0 (\partial_\mu u) - \frac{m-2}{6} g^{\mu\nu} r_0^{-m} E_0 (\partial_\mu \partial_\nu u) \\ &+ \frac{1}{2} \beta_\mu r_0^{-m} E_0 v^\mu - \frac{1}{2} r_0^{-m} E_0 (\partial_\mu v^\mu) - \frac{1}{4m} g_{\mu\nu} \sum_{\ell=0}^{m-1} r_0^{-\ell-1} r_1^{\ell-m} E_0 v^\mu E_1 v^\nu \\ &+ \frac{1}{2m} \sum_{\ell=0}^{m-1} (m-2\ell) r_0^{-\ell-1} r_1^{\ell-m} E_0 v^\mu E_1 (\partial_\mu u) + g^{\mu\nu} \sum_{\ell=0}^{m-1} \left[\frac{m-2}{6} - \frac{\ell(m-\ell-1)}{2m} \right] r_0^{-\ell-1} r_1^{\ell-m} E_0 (\partial_\mu u) E_1 (\partial_\nu u). \end{aligned}$$

Notice that:

$$g_{\mu\nu} \sum_{\ell=0}^{m-1} r_0^{-\ell-1} r_1^{\ell-m} \operatorname{tr}(E_0 v^\mu E_1 v^\nu) = g_{\mu\nu} \sum_{\ell=0}^{m-1} r_0^{-\ell-1} r_1^{\ell-m} \operatorname{tr}(E_1 v^\nu E_0 v^\mu) = g_{\mu\nu} \sum_{\ell=0}^{m-1} r_1^{-\ell-1} r_0^{\ell-m} \operatorname{tr}(E_1 v^\mu E_0 v^\nu)$$

if we change ℓ to $m - \ell - 1$ in the summation and we use the symmetry of $g_{\mu\nu}$. Then, to show that the spectral functions F reduce to the ones above, one has to use the symmetry $r_0 \leftrightarrow r_1$ for some terms. A direct computation gives

$$\begin{aligned} F_w(r_0, r_0) &= r_0^{-m}, \\ F_{\partial u}^\mu(r_0, r_0) &= \frac{m-2}{6} [\frac{1}{2} \alpha^\mu - \beta^\mu] r_0^{-m}, \\ F_{\partial \partial u}(r_0, r_0) &= -\frac{m-2}{6} r_0^{-m}, \\ F_{v, \mu}(r_0, r_0) &= \frac{1}{2} \beta_\mu r_0^{-m}, \\ F_{\partial v}(r_0, r_0) &= -\frac{1}{2} r_0^{-m}, \end{aligned}$$

$$\begin{aligned} \frac{1}{2}[F_{v,v}(r_0, r_1, r_0) + F_{v,v}(r_1, r_0, r_1)] &= -\frac{1}{4m} \sum_{\ell=0}^{m-1} r_0^{-\ell-1} r_1^{\ell-m}, \\ \frac{1}{2}[F_{\partial u, \partial u}(r_0, r_1, r_0) + F_{\partial u, \partial u}(r_1, r_0, r_1)] &= \sum_{\ell=0}^{m-1} \left[\frac{m-2}{6} - \frac{\ell(m-\ell-1)}{2m} \right] r_0^{-\ell-1} r_1^{\ell-m}, \\ F_{v, \partial u}(r_0, r_1, r_0) + F_{\partial u, v}(r_1, r_0, r_1) &= \frac{1}{2m} \sum_{\ell=0}^{m-1} (m-2\ell) r_0^{-\ell-1} r_1^{\ell-m}. \end{aligned}$$

In the last expression, under the trace we have $\text{tr}(E_0(\partial_\mu u)E_1 v^\mu) = \text{tr}(E_1 v^\mu E_0(\partial_\mu u))$, and we need to sum the two functions $F_{v, \partial u}$ and $F_{\partial u, v}$ with their correct arguments.

These relations show that (2.19) reproduces [1, eq. 4.15] when $a = \mathbb{1}$.

3.2. Minimal Laplace type operators: $u = \mathbb{1}$

Starting from (2.1) when $u = \mathbb{1}$ one gets

$$P = -g^{\mu\nu} \nabla_\mu \nabla_\nu - (p^v - [\frac{1}{2}\alpha^v - \beta^v]) \nabla_v - q$$

and two simplifications occur in (2.20): all derivatives of u vanish, and the spectrum of u reduces to $\text{sp}(u) = \{1\}$, so that all spectral functions are taken at $r_i = 1$ and $E_{r_i} = \mathbb{1}$. The result is then

$$\mathcal{R}_2 = \frac{1}{2^d \pi^{d/2}} \left[\frac{1}{6} R + G_q(1, 1)q + G_{\widehat{\nabla}_p}(1, 1)\widehat{\nabla}_\mu p^\mu + G_{p,p}(1, 1, 1)p^\mu p_\mu \right].$$

A direct computation gives $G_q(1, 1) = 1$, $G_{\widehat{\nabla}_p}(1, 1) = -\frac{1}{2}$, and $G_{p,p}(1, 1, 1) = -\frac{1}{4}$, so that

$$\mathcal{R}_2 = \frac{1}{2^d \pi^{d/2}} \left[\frac{1}{6} R + q - \frac{1}{2} \widehat{\nabla}_\mu p^\mu - \frac{1}{4} p^\mu p_\mu \right]. \tag{3.1}$$

As in Proposition 2.1, we can change $\widehat{\nabla}_\mu$ to $\widehat{\nabla}'_\mu := \widehat{\nabla}_\mu + \phi_\mu$ and solve ϕ_μ in order to get $p'^\mu = 0$. From (2.4) with $u = \mathbb{1}$, one has

$$\begin{aligned} g^{\mu\nu} \widehat{\nabla}'_\mu \widehat{\nabla}'_\nu + q' &= g^{\mu\nu} (\widehat{\nabla}_\mu + \phi_\mu) (\widehat{\nabla}_\nu + \phi_\nu) + q' = g^{\mu\nu} \widehat{\nabla}_\mu \widehat{\nabla}_\nu + g^{\mu\nu} (\widehat{\nabla}_\mu \phi_\nu) + 2g^{\mu\nu} \phi_\nu \widehat{\nabla}_\mu + g^{\mu\nu} \phi_\mu \phi_\nu + q' \\ &=: g^{\mu\nu} \widehat{\nabla}_\mu \widehat{\nabla}_\nu + p^\mu \widehat{\nabla}_\mu + q \end{aligned}$$

with $p^\mu = 2g^{\mu\nu} \phi_\nu$ and $q = q' + g^{\mu\nu} (\widehat{\nabla}_\mu \phi_\nu) + g^{\mu\nu} \phi_\mu \phi_\nu$. This is solved for $\phi_\mu = \frac{1}{2} g_{\mu\nu} p^\nu$ and implies $q' = q - \frac{1}{2} \widehat{\nabla}_\mu p^\mu - \frac{1}{4} p^\mu p_\mu$. Injected into (3.1), this gives $\mathcal{R}_2 = \frac{1}{2^d \pi^{d/2}} \left[\frac{1}{6} R + q' \right]$ as in [23, Theorem 3.3.1].

3.3. Conformal like transformed Laplacian

Let us consider a positive invertible element $k \in \Gamma(\text{End}(V))$, a covariant derivative ∇_μ on V and $\Delta = -g^{\mu\nu} \widehat{\nabla}_\mu \widehat{\nabla}_\nu$ be its associated Laplacian. Motivated by the conformal deformations worked out in [6–8], we consider the operator

$$P := k \Delta k = -g^{\mu\nu} k^2 \widehat{\nabla}_\mu \widehat{\nabla}_\nu - 2g^{\mu\nu} k (\widehat{\nabla}_\mu k) \widehat{\nabla}_\nu + k(\Delta k).$$

Actually, it is worthwhile to quote that the conformal change of metrics on the noncommutative tori is not as straightforward as copying the notation of the commutative tori since deep theories in operator algebras are involved. However, the result obtained in this section for the operator P will be used to compute \mathcal{R}_2 for the noncommutative tori in Section 5.

The operator P can be written as in (2.4) with

$$u = k^2, \quad p^v = g^{\mu\nu} k (\widehat{\nabla}_\mu k) - g^{\mu\nu} (\widehat{\nabla}_\mu k) k, \quad q = -k(\Delta k).$$

Application of Theorem 2.4 gives

$$\begin{aligned} \mathcal{R}_2^{k\Delta k} &= \frac{1}{2^d \pi^{d/2}} \left[\frac{1}{6} R r_0^{-d/2+1} E_{r_0} \right. \\ &\quad \left. + F_{\Delta k}^{k\Delta k}(r_0, r_1) E_{r_0}(\Delta k) E_{r_1} + F_{\widehat{\nabla}_k \widehat{\nabla}_k}^{k\Delta k}(r_0, r_1, r_2) g^{\mu\nu} E_{r_0}(\nabla_\mu k) E_{r_1}(\nabla_\nu k) E_{r_2} \right] \end{aligned} \tag{3.2}$$

with

$$F_{\Delta k}^{k\Delta k}(r_0, r_1) = -\sqrt{r_0} G_q(r_0, r_1) - (\sqrt{r_0} + \sqrt{r_1}) G_{\widehat{\nabla}_u}(r_0, r_1) - (\sqrt{r_0} - \sqrt{r_1}) G_{\widehat{\nabla}_p}(r_0, r_1), \tag{3.3}$$

$$\begin{aligned} F_{\widehat{\nabla}_k \widehat{\nabla}_k}^{k\Delta k}(r_0, r_1, r_2) &= 2G_{\widehat{\nabla}_u}(r_0, r_2) \\ &\quad + (\sqrt{r_0} + \sqrt{r_1})(\sqrt{r_1} + \sqrt{r_2}) G_{\widehat{\nabla}_u, \widehat{\nabla}_u}(r_0, r_1, r_2) \\ &\quad + (\sqrt{r_0} - \sqrt{r_1})(\sqrt{r_1} + \sqrt{r_2}) G_{p, \widehat{\nabla}_u}(r_0, r_1, r_2) \\ &\quad + (\sqrt{r_0} + \sqrt{r_1})(\sqrt{r_1} - \sqrt{r_2}) G_{\widehat{\nabla}_u, p}(r_0, r_1, r_2) \\ &\quad + (\sqrt{r_0} - \sqrt{r_1})(\sqrt{r_1} - \sqrt{r_2}) G_{p, p}(r_0, r_1, r_2). \end{aligned} \tag{3.4}$$

Using Proposition 2.7, one has

$$F_{\Delta k}^{k\Delta k}(r_0, r_1) = -\frac{\sqrt{r_0 r_1}(\sqrt{r_0} + \sqrt{r_1})[G_q(r_0, r_1) + 2G_{\widehat{\nu}\widehat{\nu}u}(r_0, r_1)]}{r_0 + r_1}.$$

4. Details of the computations

In this section we give some details on the computations to establish Theorems 2.3 and 2.4. These computations can be done by hand but the reader can also follow [5].

The equation of (2.19) requires to compute the terms in the sum (2.16), which itself requires to compute the arguments $B_1 \otimes \dots \otimes B_k$ and the operators $T_{k,p}$.

For $r = 2$, the list of arguments has been evaluated in [1, Section 4.1] (starting with the expression (1.1) of P) as well as their contractions with the tensor $G(g)_{\mu_1 \dots \mu_{2p}}$. We make use of these results below. Then the computation of the operators $T_{k,p}$ reduces to the computation of the universal spectral functions $I_{d/2+p,k}$. As noticed in [1], only the values $k = 1, 2, 3, 4$ have to be considered.

Below is the list of the evaluation of these arguments in the corresponding operators $T_{k,p}$, where the following functional calculus rule (and its obvious generalizations) is used

$$f(r_0, r_1, \dots, r_k) E_{r_0} u^{n_0} E_{r_1} u^{n_1} E_{r_2} \dots u^{n_{k-1}} E_{r_k} = r_0^{n_0+n_1+\dots+n_{k-1}} f(r_0, r_0, \dots, r_0) E_{r_0},$$

where summations over r_i in the spectrum of u are omitted. In the following, the symbol \rightsquigarrow is used to symbolize this evaluation.

For $k = 1$, there is only one argument:

$$w \rightsquigarrow I_{d/2,1}(r_0, r_1) E_{r_0} w E_{r_1}.$$

For $k = 2$, one has:

$$\begin{aligned} & -\frac{1}{2} g^{\mu\nu} g_{\rho\sigma} (\partial_\mu \partial_\nu g^{\rho\sigma}) u \otimes u \rightsquigarrow -\frac{1}{2} g^{\mu\nu} g_{\rho\sigma} (\partial_\mu \partial_\nu g^{\rho\sigma}) r_0^2 I_{d/2+1,2}(r_0, r_0, r_0) E_{r_0}, \\ & -g^{\mu\nu} g_{\rho\sigma} (\partial_\nu g^{\rho\sigma}) u \otimes \partial_\mu u \rightsquigarrow -g^{\mu\nu} g_{\rho\sigma} (\partial_\nu g^{\rho\sigma}) r_0 I_{d/2+1,2}(r_0, r_0, r_1) E_{r_0} (\partial_\mu u) E_{r_1}, \\ & \quad -\frac{d}{2} g^{\mu\nu} u \otimes \partial_\mu \partial_\nu u \rightsquigarrow -\frac{d}{2} g^{\mu\nu} r_0 I_{d/2+1,2}(r_0, r_0, r_1) E_{r_0} (\partial_\mu \partial_\nu u) E_{r_1}, \\ & -\frac{1}{2} g_{\rho\sigma} (\partial_\mu g^{\rho\sigma}) v^\mu \otimes u \rightsquigarrow -\frac{1}{2} g_{\rho\sigma} (\partial_\mu g^{\rho\sigma}) r_1 I_{d/2+1,2}(r_0, r_1, r_1) E_{r_0} v^\mu E_{r_1}, \\ & \quad -\frac{d}{2} v^\mu \otimes \partial_\mu u \rightsquigarrow -\frac{d}{2} I_{d/2+1,2}(r_0, r_1, r_2) E_{r_0} v^\mu E_{r_1} (\partial_\mu u) E_{r_2}, \\ & -\frac{1}{2} g_{\mu\nu} v^\mu \otimes v^\nu \rightsquigarrow -\frac{1}{2} g_{\mu\nu} I_{d/2+1,2}(r_0, r_1, r_2) E_{r_0} v^\mu E_{r_1} v^\nu E_{r_2}, \\ & \quad -u \otimes \partial_\mu v^\mu \rightsquigarrow -r_0 I_{d/2+1,2}(r_0, r_0, r_1) E_{r_0} (\partial_\mu v^\mu) E_{r_1}. \end{aligned}$$

For $k = 3$, one has:

$$\begin{aligned} & [g^{\mu\nu} g_{\rho\sigma} (\partial_\mu \partial_\nu g^{\rho\sigma}) + 2(\partial_\mu \partial_\nu g^{\mu\nu}) + g_{\rho\sigma} (\partial_\mu g^{\mu\nu})(\partial_\nu g^{\rho\sigma}) + 2g_{\rho\sigma} (\partial_\mu g^{\nu\rho})(\partial_\nu g^{\mu\sigma}) \\ & \quad + \frac{1}{2} g^{\mu\nu} g_{\rho\sigma} g_{\alpha\beta} (\partial_\mu g^{\rho\sigma})(\partial_\nu g^{\alpha\beta}) + g^{\mu\nu} g_{\rho\sigma} g_{\alpha\beta} (\partial_\mu g^{\rho\alpha})(\partial_\nu g^{\sigma\beta})] u \otimes u \otimes u \\ & \rightsquigarrow [g^{\mu\nu} g_{\rho\sigma} (\partial_\mu \partial_\nu g^{\rho\sigma}) + 2(\partial_\mu \partial_\nu g^{\mu\nu}) + g_{\rho\sigma} (\partial_\mu g^{\mu\nu})(\partial_\nu g^{\rho\sigma}) + 2g_{\rho\sigma} (\partial_\mu g^{\nu\rho})(\partial_\nu g^{\mu\sigma}) \\ & \quad + \frac{1}{2} g^{\mu\nu} g_{\rho\sigma} g_{\alpha\beta} (\partial_\mu g^{\rho\sigma})(\partial_\nu g^{\alpha\beta}) + g^{\mu\nu} g_{\rho\sigma} g_{\alpha\beta} (\partial_\mu g^{\rho\alpha})(\partial_\nu g^{\sigma\beta})] r_0^3 I_{d/2+2,3}(r_0, r_0, r_0) E_{r_0}, \\ & (d+6)[\frac{1}{2} g^{\mu\nu} g_{\rho\sigma} (\partial_\nu g^{\rho\sigma}) + (\partial_\nu g^{\mu\nu})] u \otimes u \otimes \partial_\mu u \\ & \rightsquigarrow (d+6)[\frac{1}{2} g^{\mu\nu} g_{\rho\sigma} (\partial_\nu g^{\rho\sigma}) + (\partial_\nu g^{\mu\nu})] r_0^2 I_{d/2+2,3}(r_0, r_0, r_0, r_1) E_{r_0} (\partial_\mu u) E_{r_1}, \\ & [\frac{d+4}{2} g^{\mu\nu} g_{\rho\sigma} (\partial_\nu g^{\rho\sigma}) + 2(\partial_\nu g^{\mu\nu})] u \otimes \partial_\mu u \otimes u \\ & \rightsquigarrow [\frac{d+4}{2} g^{\mu\nu} g_{\rho\sigma} (\partial_\nu g^{\rho\sigma}) + 2(\partial_\nu g^{\mu\nu})] r_0 r_1 I_{d/2+2,3}(r_0, r_0, r_1, r_1) E_{r_0} (\partial_\mu u) E_{r_1}, \\ & \frac{(d+2)^2}{2} g^{\mu\nu} u \otimes \partial_\mu u \otimes \partial_\nu u \rightsquigarrow \frac{(d+2)^2}{2} g^{\mu\nu} r_0 I_{d/2+2,3}(r_0, r_0, r_1, r_2) E_{r_0} (\partial_\mu u) E_{r_1} (\partial_\nu u) E_{r_2}, \\ & (d+2) g^{\mu\nu} u \otimes u \otimes \partial_\mu \partial_\nu u \rightsquigarrow (d+2) g^{\mu\nu} r_0^2 I_{d/2+2,3}(r_0, r_0, r_0, r_1) E_{r_0} (\partial_\mu \partial_\nu u) E_{r_1}, \\ & [\frac{1}{2} g_{\rho\sigma} (\partial_\mu g^{\rho\sigma}) + g_{\mu\nu} (\partial_\rho g^{\rho\nu})] (v^\mu \otimes u \otimes u + u \otimes v^\mu \otimes u + u \otimes u \otimes v^\mu) \\ & \rightsquigarrow [\frac{1}{2} g_{\rho\sigma} (\partial_\mu g^{\rho\sigma}) + g_{\mu\nu} (\partial_\rho g^{\rho\nu})] [r_1^2 I_{d/2+2,3}(r_0, r_1, r_1, r_1) \\ & \quad + r_0 r_1 I_{d/2+2,3}(r_0, r_0, r_1, r_1) + r_0^2 I_{d/2+2,3}(r_0, r_0, r_0, r_1)] E_{r_0} v^\mu E_{r_1}, \\ & \frac{d+2}{2} (v^\mu \otimes u \otimes \partial_\mu u) \rightsquigarrow \frac{d+2}{2} r_1 I_{d/2+2,3}(r_0, r_1, r_1, r_2) E_{r_0} v^\mu E_{r_1} (\partial_\mu u) E_{r_2}, \\ & \frac{d+2}{2} (u \otimes \partial_\mu u \otimes v^\mu) \rightsquigarrow \frac{d+2}{2} r_0 I_{d/2+2,3}(r_0, r_0, r_1, r_2) E_{r_0} (\partial_\mu u) E_{r_1} v^\mu E_{r_2}, \\ & \frac{d+2}{2} (u \otimes v^\mu \otimes \partial_\mu u) \rightsquigarrow \frac{d+2}{2} r_0 I_{d/2+2,3}(r_0, r_0, r_1, r_2) E_{r_0} v^\mu E_{r_1} (\partial_\mu u) E_{r_2}. \end{aligned}$$

For $k = 4$, one has:

$$\begin{aligned}
 & 3 \left[-\frac{1}{2} g^{\mu\nu} g_{\rho\sigma} g_{\alpha\beta} (\partial_\mu g^{\rho\sigma}) (\partial_\nu g^{\alpha\beta}) - g^{\mu\nu} g_{\rho\sigma} g_{\alpha\beta} (\partial_\mu g^{\rho\alpha}) (\partial_\nu g^{\sigma\beta}) - 2g_{\rho\sigma} (\partial_\mu g^{\mu\nu}) (\partial_\nu g^{\rho\sigma}) \right. \\
 & \quad \left. - 2g_{\rho\sigma} (\partial_\mu g^{\mu\rho}) (\partial_\nu g^{\nu\sigma}) - 2g_{\rho\sigma} (\partial_\mu g^{\nu\rho}) (\partial_\nu g^{\mu\sigma}) \right] u \otimes u \otimes u \otimes u \\
 & \rightsquigarrow 3 \left[-\frac{1}{2} g^{\mu\nu} g_{\rho\sigma} g_{\alpha\beta} (\partial_\mu g^{\rho\sigma}) (\partial_\nu g^{\alpha\beta}) - g^{\mu\nu} g_{\rho\sigma} g_{\alpha\beta} (\partial_\mu g^{\rho\alpha}) (\partial_\nu g^{\sigma\beta}) - 2g_{\rho\sigma} (\partial_\mu g^{\mu\nu}) (\partial_\nu g^{\rho\sigma}) \right. \\
 & \quad \left. - 2g_{\rho\sigma} (\partial_\mu g^{\mu\rho}) (\partial_\nu g^{\nu\sigma}) - 2g_{\rho\sigma} (\partial_\mu g^{\nu\rho}) (\partial_\nu g^{\mu\sigma}) \right] r_0^4 I_{d/2+3,4}(r_0, r_0, r_0, r_0) E_{r_0}, \\
 & - (d+4) \left[\frac{1}{2} g^{\mu\nu} g_{\rho\sigma} (\partial_\nu g^{\rho\sigma}) + (\partial_\nu g^{\mu\nu}) \right] [3u \otimes u \otimes u \otimes \partial_\mu u \\
 & \quad + 2u \otimes u \otimes \partial_\mu u \otimes u + u \otimes \partial_\mu u \otimes u \otimes u] \\
 & \rightsquigarrow -(d+4) \left[\frac{1}{2} g^{\mu\nu} g_{\rho\sigma} (\partial_\nu g^{\rho\sigma}) + (\partial_\nu g^{\mu\nu}) \right] [3r_0^3 I_{d/2+3,4}(r_0, r_0, r_0, r_1) \\
 & \quad + 2r_0^2 r_1 I_{d/2+3,4}(r_0, r_0, r_0, r_1) + r_0 r_1^2 I_{d/2+3,4}(r_0, r_0, r_1, r_1)] E_{r_0} (\partial_\mu u) E_{r_1}, \\
 & - \frac{1}{2} (d+4) (d+2) g^{\mu\nu} (2u \otimes u \otimes \partial_\mu u \otimes \partial_\nu u + u \otimes \partial_\mu u \otimes u \otimes \partial_\nu u) \\
 & \rightsquigarrow -\frac{1}{2} (d+4) (d+2) g^{\mu\nu} [2r_0^2 I_{d/2+3,4}(r_0, r_0, r_0, r_1, r_2) + r_0 r_1 I_{d/2+3,4}(r_0, r_0, r_1, r_1, r_2)] \\
 & \quad E_{r_0} (\partial_\mu u) E_{r_1} (\partial_\nu u) E_{r_2}.
 \end{aligned}$$

The coefficient c and the spectral functions F are evaluated by collecting these terms:

$$\begin{aligned}
 c & := \frac{1}{3} (\partial_\mu \partial_\nu g^{\mu\nu}) - \frac{1}{2} g^{\mu\nu} g_{\rho\sigma} (\partial_\mu \partial_\nu g^{\rho\sigma}) + \frac{1}{48} g^{\mu\nu} g_{\rho\sigma} g_{\alpha\beta} (\partial_\mu g^{\rho\sigma}) (\partial_\nu g^{\alpha\beta}) \\
 & \quad + \frac{1}{24} g^{\mu\nu} g_{\rho\sigma} g_{\alpha\beta} (\partial_\mu g^{\rho\alpha}) (\partial_\nu g^{\sigma\beta}) - \frac{1}{12} g_{\rho\sigma} (\partial_\mu g^{\mu\nu}) (\partial_\nu g^{\rho\sigma}) \\
 & \quad + \frac{1}{12} g_{\rho\sigma} (\partial_\mu g^{\nu\rho}) (\partial_\nu g^{\mu\sigma}) - \frac{1}{4} g_{\rho\sigma} (\partial_\mu g^{\mu\rho}) (\partial_\nu g^{\nu\sigma}), \\
 F_w(r_0, r_1) & := I_{d/2,1}(r_0, r_1), \\
 F_{\partial_\nu}(r_0, r_1) & := -r_0 I_{d/2+1,2}(r_0, r_0, r_1), \\
 F_{\partial\partial u}(r_0, r_1) & := -\frac{d}{2} r_0 I_{d/2+1,2}(r_0, r_0, r_1) + (d+2) r_0^2 I_{d/2+2,3}(r_0, r_0, r_0, r_1), \\
 F_{\partial u}^\mu(r_0, r_1) & := -\alpha^\mu r_0 I_{d/2+1,2}(r_0, r_0, r_1) + (d+6) \left[\frac{1}{2} \alpha^\mu + \beta^\mu \right] r_0^2 I_{d/2+2,3}(r_0, r_0, r_0, r_1) \\
 & \quad + \left[\frac{d+4}{2} \alpha^\mu + 2\beta^\mu \right] r_0 r_1 I_{d/2+2,3}(r_0, r_0, r_1, r_1) \\
 & \quad - (d+4) \left[\frac{1}{2} \alpha^\mu + \beta^\mu \right] [3r_0^3 I_{d/2+3,4}(r_0, r_0, r_0, r_1) \\
 & \quad + 2r_0^2 r_1 I_{d/2+3,4}(r_0, r_0, r_0, r_1, r_1) + r_0 r_1^2 I_{d/2+3,4}(r_0, r_0, r_1, r_1, r_1)], \\
 F_{v,\mu}(r_0, r_1) & := -\frac{1}{2} \alpha_\mu r_1 I_{d/2+1,2}(r_0, r_1, r_1) + \left[\frac{1}{2} \alpha_\mu + \beta_\mu \right] [r_1^2 I_{d/2+2,3}(r_0, r_1, r_1, r_1) \\
 & \quad + r_0 r_1 I_{d/2+2,3}(r_0, r_0, r_1, r_1) + r_0^2 I_{d/2+2,3}(r_0, r_0, r_0, r_1)], \\
 F_{v,v}(r_0, r_1, r_2) & := -\frac{1}{2} I_{d/2+1,2}(r_0, r_1, r_2), \\
 F_{\partial u,v}(r_0, r_1, r_2) & := \frac{d+2}{2} r_0 I_{d/2+2,3}(r_0, r_0, r_1, r_2), \\
 F_{v,\partial u}(r_0, r_1, r_2) & := -\frac{d}{2} I_{d/2+1,2}(r_0, r_1, r_2) + \frac{d+2}{2} r_1 I_{d/2+2,3}(r_0, r_1, r_1, r_2) \\
 & \quad + \frac{d+2}{2} r_0 I_{d/2+2,3}(r_0, r_0, r_1, r_2), \\
 F_{\partial u,\partial u}(r_0, r_1, r_2) & := \frac{(d+2)^2}{2} r_0 I_{d/2+2,3}(r_0, r_0, r_1, r_2) \\
 & \quad - \frac{(d+4)(d+2)}{2} [2r_0^2 I_{d/2+3,4}(r_0, r_0, r_0, r_1, r_2) + r_0 r_1 I_{d/2+3,4}(r_0, r_0, r_1, r_1, r_2)]. \tag{4.1}
 \end{aligned}$$

As in [1, Section 4.3], the strategy to compute (2.20) could be to make the change of variables $(u, v^\mu, w) \mapsto (u, p^\mu, q)$. Here we use another strategy which simplifies the computation since it is based on (2.8), (2.9), and (2.10).

Indeed, as already noticed, one can apply verbatim the computation of the arguments and their contractions with ∂_μ replaced by ∇_μ , and at the same time, using $p^\mu + g^{\mu\nu}(\nabla_\nu u) - [\frac{1}{2}\alpha^\mu - \beta^\mu]u$ in place of v^μ , and q in place of w . So, (2.19) can be replaced by

$$\begin{aligned}
 & c r_0^{-d/2+1} E_{r_0} + F_{\partial u}^\mu(r_0, r_1) E_{r_0} (\nabla_\mu u) E_{r_1} + g^{\mu\nu} F_{\partial\partial u}(r_0, r_1) E_{r_0} (\nabla_\mu \nabla_\nu u) E_{r_1} \\
 & + g^{\mu\nu} F_{\partial u,\partial u}(r_0, r_1, r_2) E_{r_0} (\nabla_\mu u) E_{r_1} (\nabla_\nu u) E_{r_2} + F_w(r_0, r_1) E_{r_0} w E_{r_1} + F_{v,\mu}(r_0, r_1) E_{r_0} v^\mu E_{r_1} \\
 & + F_{v,\partial u}(r_0, r_1, r_2) E_{r_0} v^\mu E_{r_1} (\nabla_\mu u) E_{r_2} + F_{\partial u,v}(r_0, r_1, r_2) a E_{r_0} (\nabla_\mu u) E_{r_1} v^\mu E_{r_2} \\
 & + g_{\mu\nu} F_{v,v}(r_0, r_1, r_2) E_{r_0} v^\mu E_{r_1} v^\nu E_{r_2} + F_{\partial v}(r_0, r_1) E_{r_0} (\nabla_\mu v^\mu) E_{r_1}.
 \end{aligned}$$

The next step is to replace ∇_μ by $\widehat{\nabla}_\mu$. We use (2.2), (2.3) and

$$\begin{aligned}
 \nabla_\mu v^\mu & = \nabla_\mu p^\mu + (\partial_\mu g^{\mu\nu}) (\nabla_\nu u) + g^{\mu\nu} (\nabla_\mu \nabla_\nu u) - \left[\frac{1}{2} \partial_\mu \alpha^\mu - \partial_\mu \beta^\mu \right] u - \left[\frac{1}{2} \alpha^\mu - \beta^\mu \right] (\nabla_\mu u) \\
 & = \nabla_\mu p^\mu + g^{\mu\nu} (\widehat{\nabla}_\mu \widehat{\nabla}_\nu u) + \beta^\mu (\widehat{\nabla}_\mu u) - \left[\frac{1}{2} \partial_\mu \alpha^\mu - \partial_\mu \beta^\mu \right] u \\
 & = \widehat{\nabla}_\mu p^\mu + \frac{1}{2} \alpha_\mu p^\mu + g^{\mu\nu} (\widehat{\nabla}_\mu \widehat{\nabla}_\nu u) + \beta^\mu (\widehat{\nabla}_\mu u) - \left[\frac{1}{2} \partial_\mu \alpha^\mu - \partial_\mu \beta^\mu \right] u.
 \end{aligned}$$

This leads to a new expression containing the terms in (2.20), with the functions given in the list after Theorem 2.4, to which we have to add the two terms $G_{\widehat{\nabla}u}^\mu(r_0, r_1)E_{r_0}(\widehat{\nabla}_\mu u)E_{r_1} + G_{p,\mu}(r_0, r_1)E_{r_0}p^\mu E_{r_1}$, with

$$\begin{aligned} G_{\widehat{\nabla}u}^\mu(r_0, r_1) &= F_{\partial u}^\mu(r_0, r_1) + g^{\mu\nu}F_{v,\nu}(r_0, r_1) + \beta^\mu F_{\partial v}(r_0, r_1) \\ &\quad + [\tfrac{1}{2}\alpha^\mu - \beta^\mu][F_{\partial\partial u}(r_0, r_1) - r_0F_{v,\partial u}(r_0, r_0, r_1) - r_1F_{\partial u,v}(r_0, r_1, r_1) \\ &\quad \quad - r_0F_{v,v}(r_0, r_0, r_1) - r_1F_{v,v}(r_0, r_1, r_1)], \\ G_{p,\mu}(r_0, r_1) &= F_{v,\mu}(r_0, r_1) + \tfrac{1}{2}\alpha_\mu F_{\partial v}(r_0, r_1) \\ &\quad - [\tfrac{1}{2}\alpha_\mu - \beta_\mu][r_0F_{v,v}(r_0, r_0, r_1) + r_1F_{v,v}(r_0, r_1, r_1)]. \end{aligned}$$

A direct computation performed using the expressions of the spectral functions F in terms of $I_{d/2,1}$ shows that

$$G_{\widehat{\nabla}u}^\mu(r_0, r_1) = G_{p,\mu}(r_0, r_1) = 0,$$

and the following symmetries

$$G_q(r_0, r_1) = G_q(r_1, r_0), \quad G_{\widehat{\nabla}\widehat{\nabla}u}(r_0, r_1) = G_{\widehat{\nabla}\widehat{\nabla}u}(r_1, r_0), \quad G_{\widehat{\nabla}u,\widehat{\nabla}u}(r_0, r_1, r_2) = G_{\widehat{\nabla}u,\widehat{\nabla}u}(r_2, r_1, r_0),$$

so that, using Proposition 2.7, one gets

$$G_{\widehat{\nabla}u,p}(r_0, r_1, r_2) = -G_{p,\widehat{\nabla}u}(r_2, r_1, r_0), \quad G_{p,p}(r_0, r_1, r_2) = G_{p,p}(r_2, r_1, r_0).$$

The coefficient in front of $r_0^{-d/2+1}E_{r_0}$ is

$$\tfrac{1}{6}R = c - \tfrac{1}{4}\alpha^\mu\beta_\mu + \tfrac{1}{2}\beta^\mu\beta_\mu + \tfrac{1}{4}\partial_\mu\alpha^\mu - \tfrac{1}{2}\partial_\mu\beta^\mu - \tfrac{1}{16}\alpha_\mu\alpha^\mu + \tfrac{1}{4}\alpha_\mu\beta^\mu - \tfrac{1}{4}\beta_\mu\beta^\mu,$$

where R is the scalar curvature of the metric g .

The spectral functions G can be written in terms of log functions for $d = 2$ (see [1, Cor. 3.3]), as Laurent polynomials for $d \geq 4$ even (see [1, Prop. 3.5]), and in terms of square roots of r_i for d odd (see [1, Prop. 3.4]). This completes the proof of Corollaries 2.8, 2.9, and 2.11.

5. Applications to the noncommutative torus

In this section, we first apply Theorem 2.4 to the noncommutative 2-torus at rational values of the deformation parameter θ , for which it is known that we get a geometrical description in terms of sections of a fiber bundle. Some computations of $a_2(a, P)$ for specific operators P have been performed at irrational values of θ to determine the so-called scalar curvature (our \mathcal{R}_2) [6–11,13–18].

We now show that we can apply our general result at rational values of θ and get the same expressions for the scalar curvature \mathcal{R}_2 which appears to be written in terms of θ -independent spectral functions. In particular, its expression is the same for rational and irrational θ .

Let $\Theta \in M_d(\mathbb{R})$ be a skew-symmetric real matrix. The noncommutative d -dimensional torus $C(\mathbb{T}_\Theta^d)$ is defined as the universal unital C^* -algebra generated by unitaries $U_k, k = 1, \dots, d$, satisfying the relations

$$U_k U_\ell = e^{2i\pi\Theta_{k,\ell}} U_\ell U_k. \tag{5.1}$$

This C^* -algebra contains, as a dense sub-algebra, the space of smooth elements for the natural action of the d -dimensional torus \mathbb{T}^d on $C(\mathbb{T}_\Theta^d)$. This sub-algebra is described as elements in $C(\mathbb{T}_\Theta^d)$ with an expansion

$$a = \sum_{(k_i) \in \mathbb{Z}^d} a_{k_1, \dots, k_d} U_1^{k_1} \dots U_d^{k_d}$$

where the sequence (a_{k_1, \dots, k_d}) belongs to the Schwartz space $S(\mathbb{Z}^d)$. We denote by $C^\infty(\mathbb{T}_\Theta^d)$ this algebra. The C^* -algebra $C(\mathbb{T}_\Theta^d)$ has a unique normalized faithful positive trace \mathfrak{t} whose restriction on smooth elements is given by

$$\mathfrak{t} \left(\sum_{(k_i) \in \mathbb{Z}^d} a_{k_1, \dots, k_d} U_1^{k_1} \dots U_d^{k_d} \right) := a_{0, \dots, 0}. \tag{5.2}$$

This trace satisfies $\mathfrak{t}(\mathbf{1}) = 1$ where $\mathbf{1}$ is the unit element of $C(\mathbb{T}_\Theta^d)$. The smooth algebra $C^\infty(\mathbb{T}_\Theta^d)$ has d canonical derivations $\delta_\mu, \mu = 1, \dots, d$, defined on the generators by

$$\delta_\mu(U_k) := \delta_{\mu,k} iU_k. \tag{5.3}$$

For any $a \in C(\mathbb{T}_\Theta^d)$, one has $\delta_\mu(a^*) = (\delta_\mu a)^*$ (real derivations).

Denote by \mathcal{H} the Hilbert space of the GNS representation of $C(\mathbb{T}_\Theta^d)$ defined by \mathfrak{t} . Each derivation δ_μ defines a unbounded operator on \mathcal{H} , denoted also by δ_μ , which satisfies $\delta_\mu^\dagger = -\delta_\mu$ (here \dagger denotes the adjoint of the operator).

5.1. The geometry of the rational noncommutative tori

In the following, we consider the special case of even dimensional noncommutative tori, $d = 2m$, with

$$\Theta = \begin{pmatrix} \theta_1 \chi & \cdots & 0 \\ 0 & \ddots & 0 \\ 0 & \cdots & \theta_m \chi \end{pmatrix}, \text{ where } \chi := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \tag{5.4}$$

for a family of deformation parameters $\theta_1, \dots, \theta_m$. Then

$$C(\mathbb{T}_\Theta^{2m}) \simeq C(\mathbb{T}_{\theta_1}^2) \otimes \cdots \otimes C(\mathbb{T}_{\theta_m}^2).$$

When $d = 2$ and $\theta = p/q$, where p, q are relatively prime integers and $q > 0$, it is known that $C(\mathbb{T}_\theta^2) \simeq \Gamma(A_\theta)$ is isomorphic to the algebra of continuous sections of a fiber bundle A_θ in $M_q(\mathbb{C})$ algebras over a 2-torus $\mathbb{T}_{\mathbb{B}^2}^2$, as recalled in Appendix A. Similarly, for $d = 4$, with $\theta_1 = p_1/q_1$ and $\theta_2 = p_2/q_2$, $C(\mathbb{T}_\Theta^4)$ is the space of sections of a fiber bundle in $M_{q_1 q_2}(\mathbb{C})$ algebras over a 4-torus $\mathbb{T}_{\mathbb{B}^2, 1}^2 \times \mathbb{T}_{\mathbb{B}^2, 2}^2$.

Moreover, in the identification $C^\infty(\mathbb{T}_\Theta^2) \simeq \Gamma^\infty(A_\theta)$, the two derivations δ_μ are the two components of the unique flat connection ∇_μ on A_θ .

This geometrical description allows to use the results of Section 2 to compute $a_2(a, P)$ for a differential operator on \mathcal{H} of the form $P = -g^{\mu\nu} u \delta_\mu \delta_\nu - [p^v + g^{\mu\nu} (\delta_\mu u) - (\frac{1}{2} \alpha^v - \beta^v) u] \delta_\nu - q$.

5.2. The noncommutative two torus

In this section, we compute the coefficient $a_2(a, P)$ on the rational noncommutative two torus for a differential operator P considered in [6,8,9,7] for the irrational noncommutative two torus. Let us introduce the following notations.

Let $\tau = \tau_1 + i\tau_2 \in \mathbb{C}$ with non zero imaginary part. We consider the constant metric g defined by

$$g^{11} = 1, \quad g^{12} = g^{21} = \Re(\tau) = \tau_1, \quad g^{22} = |\tau|^2,$$

with inverse matrix

$$g_{11} = \frac{|\tau|^2}{\Im(\tau)^2} = \frac{|\tau|^2}{\tau_2^2}, \quad g_{12} = g_{21} = -\frac{\Re(\tau)}{\Im(\tau)^2} = -\frac{\tau_1}{\tau_2^2}, \quad g_{22} = \frac{1}{\Im(\tau)^2} = \frac{1}{\tau_2^2}.$$

We will use the constant tensors

$$\epsilon^1 := 1, \quad \epsilon^2 := \tau, \quad \bar{\epsilon}^1 = 1, \quad \bar{\epsilon}^2 = \bar{\tau}, \quad h^{\mu\nu} := \bar{\epsilon}^\mu \epsilon^\nu,$$

which imply $h^{11} = 1, h^{12} = \tau, h^{21} = \bar{\tau}, h^{22} = |\tau|^2$. Then the symmetric part of $h^{\mu\nu}$ is the metric, $g^{\mu\nu} = \frac{1}{2}(h^{\mu\nu} + h^{\nu\mu})$, and $g_{\mu\nu} \bar{\epsilon}^\mu \bar{\epsilon}^\nu = 0$.

On the (GNS) Hilbert space \mathcal{H} , consider the following operators δ, δ^\dagger and the Laplacian:

$$\begin{aligned} \delta &:= \bar{\epsilon}^\mu \delta_\mu = \delta_1 + \bar{\tau} \delta_2, & \delta^\dagger &:= -\epsilon^\mu \delta_\mu = -\delta_1 - \tau \delta_2 \\ \Delta &:= \delta^\dagger \delta = -\epsilon^\mu \bar{\epsilon}^\nu \delta_\mu \delta_\nu = -h^{\mu\nu} \delta_\mu \delta_\nu = -g^{\mu\nu} \delta_\mu \delta_\nu. \end{aligned}$$

For $k \in C^\infty(\mathbb{T}_\Theta^d), k > 0$, the operator P is defined as

$$P := \begin{pmatrix} P_1 & 0 \\ 0 & P_2 \end{pmatrix} \tag{5.5}$$

with

$$\begin{aligned} P_1 &:= k \Delta k = -g^{\mu\nu} k \delta_\mu \delta_\nu k = -g^{\mu\nu} k^2 \delta_\mu \delta_\nu - 2g^{\mu\nu} k (\delta_\nu k) \delta_\mu - g^{\mu\nu} k (\delta_\mu \delta_\nu k) \\ &:= -u_1 g^{\mu\nu} \delta_\mu \delta_\nu - v_1^\mu \delta_\mu - w_1, \\ P_2 &:= \delta^\dagger k^2 \delta = -\epsilon^\nu \bar{\epsilon}^\mu \delta_\nu k^2 \delta_\mu = -\bar{\epsilon}^\mu \epsilon^\nu (\delta_\nu k^2) \delta_\mu - \bar{\epsilon}^\mu \epsilon^\nu k^2 \delta_\mu \delta_\nu = -g^{\mu\nu} k^2 \delta_\mu \delta_\nu - h^{\mu\nu} (\delta_\nu k^2) \delta_\mu \\ &:= -u_2 g^{\mu\nu} \delta_\mu \delta_\nu - v_2^\mu \delta_\mu - w_2, \end{aligned}$$

so that

$$\begin{aligned} u_1 &= k^2, & v_1^\mu &= 2g^{\mu\nu} k (\delta_\nu k), & w_1 &= -k(\Delta k), \\ u_2 &= k^2, & v_2^\mu &= h^{\mu\nu} (\delta_\nu k^2) = h^{\mu\nu} [k (\delta_\nu k) + (\delta_\nu k) k], & w_2 &= 0. \end{aligned}$$

For the forthcoming computations, since the metric g is constant, we have $c = R = \alpha^\mu = \beta^\mu = 0, \widehat{\nabla}_\mu = \nabla_\mu = \delta_\mu$ (the last equality being a property of the geometrical presentation of $C^\infty(\mathbb{T}_\Theta^d)$, as recalled above), $F_{\partial u}^\mu = F_{v, \mu} = 0$. Here $|g|^{1/2} = \det(g_{\mu\nu})^{1/2} = \tau_2^{-1}$. We can write P_1 and P_2 in the covariant form (2.1) with

$$u_1 = k^2, \quad p_1^\mu = g^{\mu\nu} [k (\delta_\nu k) - (\delta_\nu k) k], \quad q_1 = -k(\Delta k),$$

$$u_2 = k^2, \quad p_2^\mu = (h^{\mu\nu} - g^{\mu\nu})[k(\delta_\nu k) + (\delta_\nu k)k], \quad q_2 = 0.$$

Let

$$f^{\mu\nu} := \frac{1}{2}(h^{\mu\nu} - h^{\nu\mu}),$$

so that $h^{\mu\nu} = g^{\mu\nu} + f^{\mu\nu}$ and $h^{\nu\mu} = g^{\mu\nu} - f^{\mu\nu}$, and define

$$Q_g(a, b, c) := \frac{\sqrt{a}(a\sqrt{b} + 3a\sqrt{c} - \sqrt{ac} - \sqrt{abc} - 2b\sqrt{c})}{(a - b)(\sqrt{a} - \sqrt{b})(\sqrt{a} - \sqrt{c})^3},$$

$$Q_f(a, b, c) := \frac{a(\sqrt{b} + \sqrt{c})}{(a - b)(\sqrt{a} - \sqrt{b})(a - c)},$$

with the following (spectral) functions

$$F_{\Delta k}(r_0, r_1) := \frac{r_0 - r_1 - \sqrt{r_0 r_1} \ln(r_0/r_1)}{(\sqrt{r_0} - \sqrt{r_1})^3}, \tag{5.6}$$

$$F_{\partial k \partial k}^{\mu\nu}(r_0, r_1, r_2) := \frac{g^{\mu\nu}(\sqrt{r_0} + \sqrt{r_2})(\sqrt{r_0} - 2\sqrt{r_1} + \sqrt{r_2}) + f^{\mu\nu}(\sqrt{r_0} - \sqrt{r_2})^2}{(\sqrt{r_0} - \sqrt{r_1})(\sqrt{r_0} - \sqrt{r_2})^2(\sqrt{r_1} - \sqrt{r_2})} \tag{5.7}$$

$$+ [g^{\mu\nu} Q_g(r_0, r_1, r_2) - f^{\mu\nu} Q_f(r_0, r_1, r_2)] \ln(r_0/r_1)$$

$$+ [g^{\mu\nu} Q_g(r_2, r_1, r_0) - f^{\mu\nu} Q_f(r_2, r_1, r_0)] \ln(r_2/r_1).$$

As in [Appendix B](#), we use in the following result the simplified notation $\varphi(a)$ instead of $\varphi \circ L \circ S(a)$, where φ is defined in [\(1.5\)](#), while L and S are defined in [Appendix A](#). Moreover, φ and the trace t defined in [\(5.2\)](#) are related by the normalization [\(A.3\)](#).

Proposition 5.1. *For the 2-dimensional noncommutative torus at rational values of the deformation parameter θ , one has $a_2(a, P) = \varphi(a\mathcal{R}_2)$ for any $a \in C(\mathbb{T}_\theta^2)$ with*

$$\mathcal{R}_2 = \frac{1}{4\pi} [F_{\Delta k}(r_0, r_1) E_{r_0}(\Delta k) E_{r_1} + F_{\partial k \partial k}^{\mu\nu}(r_0, r_1, r_2) E_{r_0}(\delta_\mu k) E_{r_1}(\delta_\nu k) E_{r_2}]. \tag{5.8}$$

Since we are in dimension $d = 2$, the appearance of the log function in this result is expected. It is shown in [Appendix B](#) that this result coincides with a previous one in [\[8, Theorem 5.2\]](#) for the irrational noncommutative two torus.

While \mathcal{R}_2 does depend on the deformation parameter θ , and in particular if it is irrational or not, the spectral functions $F_{\Delta k}$ and $F_{\partial k \partial k}^{\mu\nu}$ do not. This universality was obtained in [\[7\]](#). Nevertheless, the fact that \mathcal{R}_2 can be written in terms of θ -independent spectral functions needs a more conceptual interpretation.

The spectrum of the differential operator P depends on the differential operators δ_μ and some multiplication operators by elements of the algebra (written here in terms of k and its derivatives $\delta_\mu k, \delta_\nu \delta_\nu k$). On the one hand, the spectrum of the closed extension of the operator δ_μ in the Hilbert space of the GNS representation consists only of eigenvalues $ik_\mu, k_\mu \in \mathbb{Z}$, associated to eigenvectors $U_1^{k_1} \cdots U_d^{k_d}$, so that it does not depend explicitly of θ . On the other hand, the computations of \mathcal{R}_2 , performed here or in [\[6,8,9,7\]](#), are based on formal manipulations of the product in the algebra, in particular they do not use the defining relations [\(5.1\)](#). This explains why these methods bypass the θ dependency and give rise to some expressions in terms of θ -independent spectral functions. Notice that for specific values of θ , for instance $\theta = 0$ (the commutative case), these expressions can be simplified. So, one has to look at [\(5.8\)](#) as a “ θ universal” expression for \mathcal{R}_2 .

Proof. One has $a_2(a, P) = a_2(a, P_1) + a_2(a, P_2)$. Denote by \mathcal{R}_2 (resp. $\mathcal{R}_2^{(1)}, \mathcal{R}_2^{(2)}$) the expressions associated to P (resp. P_1, P_2). Then one has $\mathcal{R}_2 = \mathcal{R}_2^{(1)} + \mathcal{R}_2^{(2)}$.

The operator P_1 is a conformal like transformed Laplacian, so the computation of $\mathcal{R}_2^{(1)}$ is a direct consequence of [\(3.2\)](#) in [Section 3.3](#). Here the metric is constant, so that $R = 0$, and it remains

$$\mathcal{R}_2^{(1)} := \frac{1}{4\pi} [F_{(1)\Delta k}(r_0, r_1) E_{r_0}(\Delta k) E_{r_1} + F_{(1)\partial k \partial k}^{\mu\nu}(r_0, r_1, r_2) E_{r_0}(\delta_\mu k) E_{r_1}(\delta_\nu k) E_{r_2}],$$

where, using [\(3.3\)](#) and [\(3.4\)](#),

$$F_{(1)\Delta k}(r_0, r_1) = -\sqrt{r_0} G_q(r_0, r_1) - (\sqrt{r_0} + \sqrt{r_1}) G_{\widehat{\varphi} \widehat{\varphi}_u}(r_0, r_1) - (\sqrt{r_0} - \sqrt{r_1}) G_{\widehat{\varphi}_p}(r_0, r_1),$$

$$F_{(1)\partial k \partial k}^{\mu\nu}(r_0, r_1, r_2) = 2g^{\mu\nu} G_{\widehat{\varphi} \widehat{\varphi}_u}(r_0, r_2)$$

$$+ g^{\mu\nu}(\sqrt{r_0} + \sqrt{r_1})(\sqrt{r_1} + \sqrt{r_2}) G_{\widehat{\varphi}_u, \widehat{\varphi}_u}(r_0, r_1, r_2)$$

$$+ g^{\mu\nu}(\sqrt{r_0} - \sqrt{r_1})(\sqrt{r_1} + \sqrt{r_2}) G_{p, \widehat{\varphi}_u}(r_0, r_1, r_2)$$

$$+ g^{\mu\nu}(\sqrt{r_0} + \sqrt{r_1})(\sqrt{r_1} - \sqrt{r_2}) G_{\widehat{\varphi}_u, p}(r_0, r_1, r_2)$$

$$+ g^{\mu\nu}(\sqrt{r_0} - \sqrt{r_1})(\sqrt{r_1} - \sqrt{r_2}) G_{p, p}(r_0, r_1, r_2).$$

For the operator P_2 , one applies [Theorem 2.4](#):

$$\begin{aligned} \mathcal{R}_2^{(2)} &= \frac{1}{4\pi} [(\sqrt{r_0} + \sqrt{r_1}) G_{\widehat{v}\widehat{u}}(r_0, r_1) g^{\mu\nu} E_{r_0}(\delta_\mu \delta_\nu k) E_{r_1} + 2G_{\widehat{v}\widehat{u}}(r_0, r_2) g^{\mu\nu} E_{r_0}(\delta_\mu k) E_{r_1}(\delta_\nu k) E_{r_2} \\ &\quad + (\sqrt{r_0} + \sqrt{r_1})(\sqrt{r_1} + \sqrt{r_2}) G_{\widehat{u},\widehat{v}}(r_0, r_1, r_2) g^{\mu\nu} E_{r_0}(\delta_\mu k) E_{r_1}(\delta_\nu k) E_{r_2} \\ &\quad + (\sqrt{r_0} + \sqrt{r_1})(\sqrt{r_1} + \sqrt{r_2}) G_{p,\widehat{v}}(r_0, r_1, r_2) (h^{\nu\mu} - g^{\mu\nu}) E_{r_0}(\delta_\mu k) E_{r_1}(\delta_\nu k) E_{r_2} \\ &\quad + (\sqrt{r_0} + \sqrt{r_1})(\sqrt{r_1} + \sqrt{r_2}) G_{\widehat{u},p}(r_0, r_1, r_2) (h^{\mu\nu} - g^{\mu\nu}) E_{r_0}(\delta_\mu k) E_{r_1}(\delta_\nu k) E_{r_2} \\ &\quad - (\sqrt{r_0} + \sqrt{r_1})(\sqrt{r_1} + \sqrt{r_2}) G_{p,p}(r_0, r_1, r_2) g^{\mu\nu} E_{r_0}(\delta_\mu k) E_{r_1}(\delta_\nu k) E_{r_2}] \\ &=: \frac{1}{4\pi} [F_{(2)\Delta k}(r_0, r_1) E_{r_0}(\Delta k) E_{r_1} + F_{(2)\partial k \partial k}^{\mu\nu}(r_0, r_1, r_2) E_{r_0}(\delta_\mu k) E_{r_1}(\delta_\nu k) E_{r_2}], \end{aligned}$$

with

$$\begin{aligned} F_{(2)\Delta k}(r_0, r_1) &= -(\sqrt{r_0} + \sqrt{r_1}) G_{\widehat{v}\widehat{u}}(r_0, r_1), \\ F_{(2)\partial k \partial k}^{\mu\nu}(r_0, r_1, r_2) &= 2g^{\mu\nu} G_{\widehat{v}\widehat{u}}(r_0, r_2) \\ &\quad + g^{\mu\nu} (\sqrt{r_0} + \sqrt{r_1})(\sqrt{r_1} + \sqrt{r_2}) G_{\widehat{u},\widehat{v}}(r_0, r_1, r_2) \\ &\quad + (h^{\nu\mu} - g^{\mu\nu})(\sqrt{r_0} + \sqrt{r_1})(\sqrt{r_1} + \sqrt{r_2}) G_{p,\widehat{v}}(r_0, r_1, r_2) \\ &\quad + (h^{\mu\nu} - g^{\mu\nu})(\sqrt{r_0} + \sqrt{r_1})(\sqrt{r_1} + \sqrt{r_2}) G_{\widehat{u},p}(r_0, r_1, r_2) \\ &\quad - g^{\mu\nu} (\sqrt{r_0} + \sqrt{r_1})(\sqrt{r_1} + \sqrt{r_2}) G_{p,p}(r_0, r_1, r_2). \end{aligned}$$

Then $F_{\Delta k} := F_{1,\Delta k} + F_{2,\Delta k}$ and $F_{\partial k \partial k}^{\mu\nu} := F_{1,\partial k \partial k}^{\mu\nu} + F_{2,\partial k \partial k}^{\mu\nu}$ simplify as in [\(5.6\)](#) and [\(5.7\)](#). The expression obtained for \mathcal{R}_2 shows that it belongs to $C(\mathbb{T}_\theta^2)$ and acts by left multiplication on \mathcal{H} . ■

5.3. The noncommutative four torus

Our result applies to the computation of the conformally perturbed scalar curvature on the noncommutative four torus, computed in [\[11,14\]](#). In order to do that, as in dimension 2, we perform the computation at rational value of θ as described at the end of [Appendix A](#).

The operator we consider is the one in [\[11\]](#), written as

$$\Delta_\varphi := k^2 \bar{\partial}_1 k^{-2} \partial_1 k^2 + k^2 \partial_1 k^{-2} \bar{\partial}_1 k^2 + k^2 \bar{\partial}_2 k^{-2} \partial_2 k^2 + k^2 \partial_2 k^{-2} \bar{\partial}_2 k^2$$

with (in our notations) $k^2 := e^h$, $\partial_1 := -\delta_1 + i\delta_3$, $\bar{\partial}_1 := -\delta_1 - i\delta_3$, $\partial_2 := -\delta_2 + i\delta_4$, and $\bar{\partial}_2 := -\delta_2 - i\delta_4$. Indeed, in [\[11,14\]](#), the derivations are $\widehat{\delta}_\mu = -i\delta_\mu$. This leads to

$$\begin{aligned} \Delta_\varphi &= -2 [k^2 g^{\mu\nu} \delta_\mu \delta_\nu + g^{\mu\nu} (\delta_\nu k^2) \delta_\mu + g^{\mu\nu} (\delta_\mu \delta_\nu k^2) - g^{\mu\nu} (\delta_\mu k^2) k^{-2} (\delta_\nu k^2)] \\ &=: 2P. \end{aligned}$$

The metric $g^{\mu\nu}$ is the diagonal one in [\[11\]](#), but in the following computation, we only require $g^{\mu\nu}$ to be constant.

Let us mention that in [\[11,14\]](#), the computation of the scalar curvature is done using P defined above (and not Δ_φ), since the symbol in [\[11, Lemma 3.6\]](#) is the one of P . So we will use P in the following. We get $P =: -ug^{\mu\nu} \delta_\mu \delta_\nu - v^\mu \delta_\mu - w$ with

$$u = k^2, \quad v^\mu = g^{\mu\nu} (\delta_\nu k^2), \quad w = g^{\mu\nu} (\delta_\mu \delta_\nu k^2) - g^{\mu\nu} (\delta_\mu k^2) k^{-2} (\delta_\nu k^2).$$

Since g is constant, we have as before $c = R = \alpha^\mu = \beta^\mu = 0$ and $\widehat{\nabla}_\mu = \nabla_\mu = \delta_\mu$ and this implies $p^\mu = 0$ and $q = g^{\mu\nu} (\delta_\mu \delta_\nu k^2) - g^{\mu\nu} (\delta_\mu k^2) k^{-2} (\delta_\nu k^2)$ in the covariant form [\(2.1\)](#). We then use the result of [Corollary 2.10](#) to get the conformally perturbed scalar curvature:

$$\mathcal{R}_2 = \frac{1}{2^5 \pi^2} [g^{\mu\nu} k^{-2} (\delta_\mu \delta_\nu k^2) k^{-2} - \frac{3}{2} g^{\mu\nu} k^{-2} (\delta_\mu k^2) k^{-2} (\delta_\nu k^2) k^{-2}]. \tag{5.9}$$

In [Appendix B](#) it is shown that we recover the result previously obtained in [\[11,14\]](#) for the irrational noncommutative four torus.

6. Conclusion

In this paper, we have computed in all dimensions the local section \mathcal{R}_2 of $\text{End}(V)$ defined by $a_2(a, P) = \int_M \text{tr}[a(x) \mathcal{R}_2(x)] \text{dvol}_g(x)$ for any section a of $\text{End}(V)$ for any nonminimal Laplace type operator $P = -[g^{\mu\nu} u(x) \partial_\mu \partial_\nu + v^\nu(x) \partial_\nu + w(x)]$ ([Theorems 2.3](#) and [2.4](#)). Expressions have been given for \mathcal{R}_2 in small dimensions, $d = 2, 3, 4$ ([Corollary 2.8, 2.10, and 2.11](#)) and for any even dimension $d \geq 2$ ([Corollary 2.9](#)), where, as expected from the results in [\[1\]](#), polynomials expressions can be proposed.

Despite the difficulties, $a_4(a, P)$ has been exhibited for $d = 2$ in [\[18\]](#) for the 2-dimensional noncommutative torus, leaving open the computation of \mathcal{R}_4 . We hope that our method could be used to reach \mathcal{R}_4 in any dimension, using a computer algebra system in the more general framework of an arbitrary P , like [\(1.1\)](#).

Our method still applies to more general setting than the NCT at rational values of the deformation parameter, namely to n -homogeneous C^* -algebras, which can be characterized in terms of sections of fiber bundles with fiber space $M_n(\mathbb{C})$ [\[24,25\]](#).

Acknowledgments

The authors are indebted to Valentin Zagrebnev for helpful discussions concerning some aspects of this paper and the referee for its pertinent suggestions.

Appendix A. Geometrical identification of the noncommutative torus at rational values

For $d = 2$ and $\theta := p/q$ rational, with p, q relatively prime integers with $q > 0$, it is known (see [26, Prop. 12.2], [27, Sect. 3]) that the algebra $C(\mathbb{T}_\theta^2)$ of the NCT identifies with the algebra $\Gamma(A_\theta)$ of continuous sections of a fiber bundle A_θ in $M_q(\mathbb{C})$ algebras over a 2-torus \mathbb{T}_p^2 . Let us describe this identification.

Denote by \mathbb{T}_p^2 the 2-torus given by identification of opposite sides of the square $[0, 2\pi]^2$. An element in \mathbb{T}_p^2 is written as (e^{ix}, e^{iy}) for $(x, y) \in [0, 2\pi]^2$. There is a natural action of the (abelian discrete) group $G := \mathbb{Z}_q^2$ on \mathbb{T}_p^2 : $(m, n) \cdot (e^{ix}, e^{iy}) := (e^{i(x+2\pi pm/q)}, e^{i(y+2\pi pn/q)})$.

The quotient $\mathbb{T}_B^2 := \mathbb{T}_p^2/G$ is the 2-torus constructed by identification of opposite sides of the square $[0, 2\pi/q]^2$. Indeed, there are unique $m \in \mathbb{Z}_q$ and $n \in \mathbb{Z}_q$ such that $e^{i(x+2\pi pm/q)} = e^{i(x+2\pi/q)}$ and $e^{i(y+2\pi pn/q)} = e^{i(y+2\pi/q)}$, so that $(m, 0)$ (resp. $(0, n)$) identifies (e^{ix}, e^{iy}) with $(e^{i(x+2\pi/q)}, e^{iy})$ (resp. $(e^{ix}, e^{i(y+2\pi/q)})$) in \mathbb{T}_p^2/G . The quotient map $\mathbb{T}_p^2 \rightarrow \mathbb{T}_B^2$ is a G -covering.

Let us now consider the C^* -algebra $C(\mathbb{T}_p^2, M_q(\mathbb{C})) \simeq C(\mathbb{T}_p^2) \otimes M_q(\mathbb{C})$ of matrix-valued continuous functions on \mathbb{T}_p^2 , in which the space of smooth functions $C^\infty(\mathbb{T}_p^2, M_q(\mathbb{C}))$ is a dense subalgebra. In order to describe this algebra, let us consider the two matrices

$$U_0 := \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & & & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \\ 1 & 0 & 0 & \dots & 0 \end{pmatrix}, \quad V_0 := \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & \xi_1 & 0 & \dots & 0 \\ \vdots & & \ddots & & \vdots \\ 0 & \dots & 0 & \xi_{q-2} & 0 \\ 0 & 0 & \dots & 0 & \xi_{q-1} \end{pmatrix} \quad \text{with } \xi_n := e^{i2\pi n\theta},$$

which satisfy $U_0 V_0 = e^{i2\pi\theta} V_0 U_0$, $U_0^q = V_0^q = \mathbb{1}_q$. For $(r, s) \in \mathbb{Z}_q^2$, the $U_0^r V_0^s$'s define a basis of $M_q(\mathbb{C})$ such that $\text{tr}[U_0^r V_0^s] = q \delta_{(r,s),(0,0)}$ (here tr is the trace on $M_q(\mathbb{C})$). Then $a \in C^\infty(\mathbb{T}_p^2, M_q(\mathbb{C}))$ can be decomposed as

$$a(e^{ix}, e^{iy}) = \sum_{(r,s) \in \mathbb{Z}_q^2} a_{r,s}(e^{ix}, e^{iy}) U_0^r V_0^s = \sum_{(k,\ell) \in \mathbb{Z}^2} \sum_{(r,s) \in \mathbb{Z}_q^2} a_{k,\ell,r,s} u(x)^k v(y)^\ell U_0^r V_0^s \tag{A.1}$$

where, with $u(x) := e^{ix}$ and $v(y) := e^{iy}$, the last decomposition is the Fourier series of the smooth functions $a_{r,s}$ on \mathbb{T}_p^2 . In particular, $a_{k,\ell,r,s}$ are rapidly decreasing coefficients in terms of $(k, \ell) \in \mathbb{Z}^2$.

The group G acts on $M_q(\mathbb{C})$ by

$$(m, n) \cdot A := U_0^{-n} V_0^m A V_0^{-m} U_0^n.$$

Let us consider the subalgebra $C_G(\mathbb{T}_p^2, M_q(\mathbb{C})) \subset C(\mathbb{T}_p^2, M_q(\mathbb{C}))$ of G -equivariant functions, which by definition satisfy, for any $(m, n) \in G$,

$$a(e^{i(x+2\pi pm/q)}, e^{i(y+2\pi pn/q)}) = U_0^n V_0^{-m} a(e^{ix}, e^{iy}) V_0^m U_0^{-n}.$$

In the form of (A.1), the G -equivariant elements in $C^\infty(\mathbb{T}_p^2, M_q(\mathbb{C}))$ are such that their coefficients satisfy $a_{k,\ell,r,s} e^{i2\pi(mk+n\ell)} = a_{k,\ell,r,s} e^{i2\pi(mr+ns)}$ for any $(m, n) \in G$, $(k, \ell) \in \mathbb{Z}^2$ and $(r, s) \in \mathbb{Z}_q^2$, so that $a_{k,\ell,r,s} \neq 0$ only when $mk + n\ell \equiv mr + ns \pmod q$. With $(m, n) = (1, 0)$ and $(0, 1)$ this implies $k \equiv r \pmod q$ and $\ell \equiv s \pmod q$. In (A.1), for a couple $(k, \ell) \in \mathbb{Z}^2$, there is a unique $(r, s) \in \mathbb{Z}_q^2$ for which $a_{k,\ell,r,s} \neq 0$ (r and s are the remainders of the Euclidean divisions of k and ℓ by q). Then, the only non zero coefficients $a_{k,\ell,r,s}$ depend only on $(k, \ell) \in \mathbb{Z}^2$. We denote them by $a_{k,\ell}$, and a smooth G -equivariant function $a \in C_G^\infty(\mathbb{T}_p^2, M_q(\mathbb{C}))$ is then given by the expansion

$$a = \sum_{(k,\ell) \in \mathbb{Z}^2} a_{k,\ell} (uU_0)^k (vV_0)^\ell = \sum_{(k,\ell) \in \mathbb{Z}^2} a_{k,\ell} U^k V^\ell$$

with $U := uU_0$, $V := vV_0$ satisfying $UV = e^{i2\pi\theta} VU$. Then, the C^* -algebra $C(\mathbb{T}_\theta^2)$ for $\theta = p/q$ identifies with $C_G(\mathbb{T}_p^2, M_q(\mathbb{C}))$, the C^* -completion of $C_G^\infty(\mathbb{T}_p^2, M_q(\mathbb{C}))$ in $C(\mathbb{T}_p^2, M_q(\mathbb{C}))$.

The space $C_G(\mathbb{T}_p^2, M_q(\mathbb{C}))$ identifies in a canonical way with the space $\Gamma(A_\theta)$ of continuous sections of the associated fiber bundle $A_\theta := \mathbb{T}_p^2 \times_G M_q(\mathbb{C})$ to the G -covering $\mathbb{T}_p^2 \rightarrow \mathbb{T}_B^2$. By definition, A_θ is the quotient of $\mathbb{T}_p^2 \times M_q(\mathbb{C})$ by the equivalence relation $((m, n) \cdot (e^{ix}, e^{iy}), A) \sim ((e^{ix}, e^{iy}), (m, n) \cdot A)$ for any $(m, n) \in G$. We denote by $[(e^{ix}, e^{iy}), A] \in A_\theta$ the class of $((e^{ix}, e^{iy}), A)$. We denote by $S : C_G(\mathbb{T}_p^2, M_q(\mathbb{C})) \rightarrow \Gamma(A_\theta)$ the identification, defined by $S(a)(x, y) := [(e^{ix}, e^{iy}), a(e^{ix}, e^{iy})]$.

In the GNS construction, $C_G^\infty(\mathbb{T}_p^2, M_q(\mathbb{C}))$ is dense in \mathcal{H} and is contained in the domains of the δ_μ 's. The fiber of the vector bundle V on which the differential operator P acts is then $\mathbb{C}^N \simeq M_q(\mathbb{C})$, i.e. $N = q^2$. In the present situation, all used elements in $\Gamma(\text{End}(V))$ are in fact left multiplications by elements in $C_G(\mathbb{T}_p^2, M_q(\mathbb{C})) \simeq \Gamma(A_\theta)$. For instance, the element a in (1.4) will be understood as the left multiplication by an element $a \in C_G(\mathbb{T}_p^2, M_q(\mathbb{C}))$.

For $A \in M_q(\mathbb{C})$, let $L(A)$ be the left multiplication by A on $M_q(\mathbb{C})$. Then $L(A)$ has the same spectrum as A , each eigenvalues having q times its original multiplicity. In particular, we have that

$$\text{tr}[L(A)] = q \text{tr}[A]$$

where in the LHS tr is the trace of operators on $M_q(\mathbb{C})$ and in the RHS tr is the trace on $M_q(\mathbb{C})$.

The computation of \mathcal{R}_2 uses local trivializations of sections of A_θ . Given a section $S(a) \in \Gamma(A_\theta)$ associated to $a \in C_C^\infty(\mathbb{T}_B^2, M_q(\mathbb{C}))$, we define the local section $S(a)_{\text{loc}} : (0, 2\pi/q)^2 \rightarrow M_q(\mathbb{C})$ by $S(a)_{\text{loc}}(x, y) := a(e^{ix}, e^{iy})$. Notice that the open subset $(0, 2\pi/q)^2 \subset \mathbb{T}_B^2$ is sufficient to describe the continuous section $S(a)$ via its trivialization $S(a)_{\text{loc}}$. The (local) section \mathcal{R}_2 relative to φ in (1.4) is defined by

$$\varphi(L(S(a)_{\text{loc}})) = \int_{\mathbb{T}_B^2} \text{tr}[L(S(a)_{\text{loc}}(x, y))] \text{dvol}_g(x, y) = |g|^{1/2} \int_0^{2\pi/q} dx \int_0^{2\pi/q} dy q \text{tr}[S(a)_{\text{loc}}(x, y)].$$

where we suppose here that $|g|^{1/2}$ is constant (this is the case for the situations considered in the paper). Then, one has

$$S(a)_{\text{loc}}(x, y) = \sum_{(k, \ell) \in \mathbb{Z}^2} a_{k, \ell} (u(x)U_0)^k (v(x)V_0)^\ell,$$

hence

$$\varphi(L(S(a)_{\text{loc}})) = |g|^{1/2} q \sum_{(k, \ell) \in \mathbb{Z}^2} a_{k, \ell} \int_0^{2\pi/q} dx e^{ikx} \int_0^{2\pi/q} dy e^{i\ell y} \text{tr}[U_0^k V_0^\ell].$$

The trace $\text{tr}[U_0^k V_0^\ell]$ is non-zero only when k, ℓ are multiples of q , and its value is then q , so that

$$\varphi(L(S(a)_{\text{loc}})) = |g|^{1/2} q^2 \sum_{(k, \ell) \in \mathbb{Z}^2} a_{k, \ell} \int_0^{2\pi/q} dx e^{iqkx} \int_0^{2\pi/q} dy e^{iq\ell y} = |g|^{1/2} q^2 \left(\frac{2\pi}{q}\right)^2 a_{0,0} = (2\pi)^2 |g|^{1/2} \mathfrak{t}(a). \tag{A.2}$$

Finally we get (when $|g|^{1/2}$ is constant)

$$\varphi \circ L \circ S = (2\pi)^2 |g|^{1/2} \mathfrak{t} \tag{A.3}$$

when applied to any elements in $C(\mathbb{T}_\Theta^2)$.

Consider now a 4-dimensional noncommutative torus for $\Theta = \begin{pmatrix} \theta_1 \chi & 0 \\ 0 & \theta_2 \chi \end{pmatrix}$, as in (5.4), and $\theta_i = p_i/q_i, p_i, q_i$ relatively prime integers, and $q_i > 0$.

Then $C(\mathbb{T}_\Theta^4) = C(\mathbb{T}_{\Theta_1}^2) \otimes C(\mathbb{T}_{\Theta_2}^2) = \Gamma(A_{\theta_1}) \otimes \Gamma(A_{\theta_2}) = \Gamma(A_{\theta_1} \boxtimes A_{\theta_2})$ where $A_{\theta_1} \boxtimes A_{\theta_2}$ is the external tensor product of the two vector bundles A_{θ_i} over the base 2-torus $\mathbb{T}_{B,i}^2$ defined as above. Recall that, with $\text{pr}_i : \mathbb{T}_{B,1}^2 \times \mathbb{T}_{B,2}^2 \rightarrow \mathbb{T}_{B,i}^2$ the natural projections, $A_{\theta_1} \boxtimes A_{\theta_2} := (\text{pr}_1^* A_{\theta_1}) \otimes (\text{pr}_2^* A_{\theta_2})$ where $\text{pr}_i^* A_{\theta_i}$ is the pull-back of A_{θ_i} on $\mathbb{T}_{B,1}^2 \times \mathbb{T}_{B,2}^2$. The fiber of $A_{\theta_1} \boxtimes A_{\theta_2}$ is then $M_{q_1}(\mathbb{C}) \otimes M_{q_2}(\mathbb{C}) \simeq M_{q_1 q_2}(\mathbb{C})$ and the isomorphism $\Gamma(A_{\theta_1}) \otimes \Gamma(A_{\theta_2}) \xrightarrow{\simeq} \Gamma(A_{\theta_1} \boxtimes A_{\theta_2})$ is induced by $(s_1 \otimes s_2)(x_1, x_2) := s_1(x_1) \otimes s_2(x_2)$ for any $s_i \in \Gamma(A_{\theta_i})$ and $x_i \in \mathbb{T}_{B,i}^2$. Using the same line of arguments as for the 2-dimensional case, and denoting by g_i a constant metric on $\mathbb{T}_{B,i}^2$, one gets, for any $a_i \in C(\mathbb{T}_{\Theta_i}^2)$,

$$\varphi \circ L(S(a_1) \otimes S(a_2)) = |g_1|^{1/2} |g_2|^{1/2} (q_1 q_2)^2 \left(\frac{2\pi}{q_1}\right)^2 \left(\frac{2\pi}{q_2}\right)^2 \mathfrak{t}(a_1 \otimes a_2), \tag{A.4}$$

so that

$$\varphi \circ L \circ (S \otimes S) = (2\pi)^4 |g_1|^{1/2} |g_2|^{1/2} \mathfrak{t}. \tag{A.5}$$

This procedure can be extended straightforwardly to any even dimension.

Appendix B. Comparison with previous results for noncommutative tori

We would like to compare the result (5.8) with [8, Theorem 5.2]. Some transformations are in order, since some conventions are different and the results are presented using different operators. In [8, Theorem 5.2], it is presented relative to the normalized trace \mathfrak{t} on $C(\mathbb{T}_\Theta^2)$, while our result is presented relative to $\varphi \circ L \circ S = (2\pi)^2 |g|^{1/2} \mathfrak{t}$ with here $|g|^{1/2} = \tau_2^{-1}$. If \mathcal{R}_{FK} denotes the operator of [8, Theorem 5.2], then

$$a_2(a, P) = \mathfrak{t}(a \mathcal{R}_{\text{FK}}) = \varphi(a \mathcal{R}_2) = (2\pi)^2 \tau_2^{-1} \mathfrak{t}(a \mathcal{R}_2), \quad \forall a \in C(\mathbb{T}_\Theta^2),$$

so we need to show that $\mathcal{R}_2 = \frac{\tau_2}{(2\pi)^2} \mathcal{R}_{\text{FK}}$ (strictly speaking, $a \mathcal{R}_2$ should be replaced by $L(S(a) \mathcal{R}_2)$ in $\varphi(a \mathcal{R}_2)$, see Appendix A).

Present results are given using functional calculus on the left and right multiplication operators L_u and R_u where $u = k^2$. The corresponding spectral decompositions give $L_u(a) = \sum_{r_0} r_0 E_{r_0} a$ and $R_u(a) = \sum_{r_1} r_1 a E_{r_1}$, where E_{r_i} is the projection

associated to u for the spectral value r_i . In [8], another convention is used, namely via functional calculus on the modular operator $\Delta(a) := k^{-2}ak^2$. If E_y^Δ denotes the projection of Δ associated to the spectral value y , then

$$\Delta(a) = L_{u^{-1}} \circ R_u(a) = \sum_{r_0, r_1} r_0^{-1} r_1 E_{r_0} a E_{r_1} = \sum_{r_0, y} y E_{r_0} a E_{y r_0} = \sum_y y E_y^\Delta(a)$$

where $y := r_0^{-1} r_1$ belongs to the spectrum of Δ .

Thus $E_y^\Delta(a) = \sum_{r_0} E_{r_0} a E_{y r_0}$ and is easy to check that $E_{y_0}^\Delta E_{y_1}^\Delta = \delta_{y_0, y_1} E_{y_0}^\Delta$.

The next three lemmas are technical results which permit to transform our relation (5.8) into a relation that will be compared to [8, Theorem 5.2].

For any $b_0 \otimes \dots \otimes b_p \in M_N(\mathbb{C})^{\otimes p+1}$, we denote by Δ_i the operator Δ which acts on b_i , by L_k^i the left multiplication by k acting on b_i , and by R_u^i the right multiplication by u on b_i . Notice that all these operators commute.

The first lemma transforms the functional calculus in the R_u^i 's into a functional calculus in the Δ_i 's.

Lemma B.1 (Substitution Lemma). For any function $F(r_0, r_1, \dots, r_p)$ of the eigenvalues r_i of the R_u^i 's and any $b_0 \otimes \dots \otimes b_p$, one has

$$\sum_{r_0, r_1, \dots, r_p} F(r_0, r_1, \dots, r_p) b_0 E_{r_0} b_1 E_{r_1} \dots b_p E_{r_p} = \sum_{r_0, y_1, \dots, y_p} f(r_0, y_1, \dots, y_p) b_0 E_{r_0} E_{y_1}^\Delta(b_1) \dots E_{y_p}^\Delta(b_p)$$

where $f(r_0, y_1, \dots, y_p) := F(r_0, r_0 y_1, r_0 y_1 y_2, \dots, r_0 y_1 \dots y_p)$ is a spectral function of R_u^0 and the Δ_i 's.

Using functional calculus notation, this lemma implies

$$\mathbf{m} \circ F(R_u^0, R_u^1, \dots, R_u^p) = \mathbf{m} \circ f(R_u^0, \Delta_1, \dots, \Delta_p)$$

as operators acting on elements $b_0 \otimes \dots \otimes b_p$. This result is very analog to the rearrangement lemma [28, Corollary 3.9] without the integral $\int_0^\infty du$ in [28, eq. (3.9)].

Proof. It is sufficient to show how the combinatorial aspect of the proof works for $p = 2$. One has

$$\begin{aligned} \sum_{r_0, y_1, y_2} f(r_0, y_1, y_2) b_0 E_{r_0} E_{y_1}^\Delta(b_1) E_{y_2}^\Delta(b_2) &= \sum_{r_0, y_1, y_2} F(r_0, r_0 y_1, r_0 y_1 y_2) b_0 E_{r_0} E_{y_1}^\Delta(b_1) E_{y_2}^\Delta(b_2) \\ &= \sum_{\substack{r_0, y_1, y_2 \\ z_1, z_2}} F(r_0, r_0 y_1, r_0 y_1 y_2) b_0 E_{r_0} E_{z_1} b_1 E_{y_1 z_1} E_{z_2} b_2 E_{y_2 z_2} \\ &= \sum_{r_0, y_1, y_2, z_2} F(r_0, r_0 y_1, r_0 y_1 y_2) b_0 E_{r_0} b_1 E_{y_1 r_0} E_{z_2} b_2 E_{y_2 z_2} \\ &= \sum_{r_0, r_1, y_2, z_2} F(r_0, r_1, r_1 y_2) b_0 E_{r_0} b_1 E_{r_1} E_{z_2} b_2 E_{y_2 z_2} \\ &= \sum_{r_0, r_1, y_2} F(r_0, r_1, r_1 y_2) b_0 E_{r_0} b_1 E_{r_1} b_2 E_{y_2 r_1} \\ &= \sum_{r_0, r_1, r_2} F(r_0, r_1, r_2) b_0 E_{r_0} b_1 E_{r_1} b_2 E_{r_2}. \quad \blacksquare \end{aligned}$$

Let $k = e^{h/2}$. While the arguments b_i mentioned above are $\delta_\mu k$ or Δk , they are $\delta_\mu(\ln k) = \frac{1}{2} \delta_\mu h$ and $\Delta(\ln k) = \frac{1}{2} \Delta h$ in [7]. The second lemma gives the relations between these arguments, compare with [8, Lemma 5.1].

Lemma B.2. If $g_1(y) = \frac{\sqrt{y}-1}{\ln y}$ and $g_2(y_1, y_2) = 2 \frac{\sqrt{y_1}(\sqrt{y_2}-1) \ln y_1 - (\sqrt{y_1}-1) \ln y_2}{\ln y_1 \ln y_2 (\ln y_1 + \ln y_2)}$, then

$$\begin{aligned} \delta_\mu k &= k g_1(\Delta) [\delta_\mu h] = 2k g_1(\Delta) [\delta_\mu \ln k], \\ \Delta k &= k g_1(\Delta) [\Delta h] - g^{\mu\nu} k \mathbf{m} \circ g_2(\Delta_1, \Delta_2) [(\delta_\mu h) \otimes (\delta_\nu h)] \\ &= 2k g_1(\Delta) [\Delta \ln k] - 4g^{\mu\nu} k \mathbf{m} \circ g_2(\Delta_1, \Delta_2) [(\delta_\mu \ln k) \otimes (\delta_\nu \ln k)]. \end{aligned}$$

Proof. With $g_1(y) := \frac{1}{2} \int_0^1 ds_1 y^{s_1/2} = (\sqrt{y} - 1) \ln^{-1} y$, we get

$$\begin{aligned} \delta_\mu k &= \delta_\mu e^{h/2} = \int_0^1 ds_1 e^{(1-s_1)h/2} (\delta_\mu h/2) e^{s_1 h/2} = \frac{1}{2} k \left(\int_0^1 ds_1 \Delta^{s_1/2} \right) [\delta_\mu h] \\ &= k \frac{\Delta^{1/2} - 1}{\ln \Delta} [\delta_\mu h] = 2k \frac{\Delta^{1/2} - 1}{\ln \Delta} [\delta_\mu \ln k] = k g_1(\Delta) [\delta_\mu h] = 2k g_1(\Delta) [\delta_\mu \ln k]. \end{aligned}$$

Similarly for the Laplacian,

$$\begin{aligned} \Delta k &= -g^{\mu\nu}(\delta_\mu \delta_\nu k) = -\frac{1}{2}g^{\mu\nu} \delta_\mu \left[\int_0^1 ds_1 e^{(1-s_1)h/2} (\delta_\nu h) e^{s_1 h/2} \right] \\ &= -\frac{1}{2}g^{\mu\nu} \left[\int_0^1 ds_1 e^{(1-s_1)h/2} (\delta_\mu \delta_\nu h) e^{s_1 h/2} \right] \\ &\quad - \frac{1}{4}g^{\mu\nu} \left[\int_0^1 ds_1 \int_0^{1-s_1} ds_2 e^{(1-s_1-s_2)h/2} (\delta_\mu h) e^{s_2 h/2} (\delta_\nu h) e^{s_1 h/2} \right] \\ &\quad - \frac{1}{4}g^{\mu\nu} \left[\int_0^1 ds_1 \int_0^{s_1} ds_2 e^{(1-s_1)h/2} (\delta_\nu h) e^{(s_1-s_2)h/2} (\delta_\mu h) e^{s_2 h/2} \right] \\ &= -\frac{1}{2}g^{\mu\nu} k \left[\int_0^1 ds_1 \Delta^{s_1/2} \right] (\delta_\mu \delta_\nu h) \\ &\quad - \frac{1}{4}g^{\mu\nu} k \mathbf{m} \circ \left[\int_0^1 ds_1 \int_0^{1-s_1} ds_2 \Delta_1^{(s_1+s_2)/2} \Delta_2^{s_1/2} \right] [(\delta_\mu h) \otimes (\delta_\nu h)] \\ &\quad - \frac{1}{4}g^{\mu\nu} k \mathbf{m} \circ \left[\int_0^1 ds_1 \int_0^{s_1} ds_2 \Delta_1^{s_1/2} \Delta_2^{s_2/2} \right] [(\delta_\mu h) \otimes (\delta_\nu h)] \\ &= k g_1(\Delta) [\Delta h] - g^{\mu\nu} k \mathbf{m} \circ g_2(\Delta_1, \Delta_2) [(\delta_\mu h) \otimes (\delta_\nu h)] \end{aligned}$$

with

$$\begin{aligned} g_2(y_1, y_2) &:= \frac{1}{4} \int_0^1 ds_1 \int_0^{1-s_1} ds_2 y_1^{(s_1+s_2)/2} y_2^{s_1/2} + \frac{1}{4} \int_0^1 ds_1 \int_0^{s_1} ds_2 y_1^{s_1/2} y_2^{s_2/2} \\ &= 2 \frac{\sqrt{y_1}(\sqrt{y_2}-1) \ln y_1 - (\sqrt{y_1}-1) \ln y_2}{\ln y_1 \ln y_2 (\ln y_1 + \ln y_2)}. \quad \blacksquare \end{aligned}$$

The third lemma gives (technical) functional relations which allow a change of arguments inside our operators. Denote by $\mathbf{m}_{12}[b_0 \otimes b_1 \otimes b_2] := b_0 \otimes b_1 b_2$ the partial multiplication.

Lemma B.3. For any operators like $f_1(R_u^0, \Delta_1), f_2(R_u^0, \Delta_1, \Delta_2), g_1(\Delta_1)$, and $g_2(\Delta_1, \Delta_2)$, one has

$$\begin{aligned} \mathbf{m} \circ f_1(R_u^0, \Delta_1) \circ L_k^1 \circ g_1(\Delta_1) &= \mathbf{m} \circ R_k^0 \circ f_1(R_u^0, \Delta_1) \circ g_1(\Delta_1), \\ \mathbf{m} \circ f_1(R_u^0, \Delta_1) \circ \mathbf{m}_{12} \circ L_k^1 \circ g_2(\Delta_1, \Delta_2) &= \mathbf{m} \circ R_k^0 \circ f_1(R_u^0, \Delta_1 \Delta_2) \circ g_2(\Delta_1, \Delta_2), \\ \mathbf{m} \circ f_2(R_u^0, \Delta_1, \Delta_2) \circ L_k^1 \circ L_k^2 \circ g_2(\Delta_1, \Delta_2) &= \mathbf{m} \circ R_u^0 \circ \Delta_1^{1/2} \circ f_2(R_u^0, \Delta_1, \Delta_2) \circ g_2(\Delta_1, \Delta_2). \end{aligned}$$

Thus the operators on the LHS are respectively associated, modulo the multiplication operator \mathbf{m} , to operators defined by the spectral functions

$$\sqrt{r_0} f_1(r_0, y_1) g_1(y_1), \quad \sqrt{r_0} f_1(r_0, y_1 y_2) g_2(y_1, y_2), \quad r_0 \sqrt{y_1} f_2(r_0, y_1, y_2) g_2(y_1, y_2),$$

where y_1, y_2 belong to the spectrum of Δ and r_0 to the spectrum of u .

Proof. For the first relation, we compute the LHS on $b_0 \otimes b_1$ using spectral decomposition:

$$\begin{aligned} \mathbf{m} \circ f_1(R_u^0, \Delta_1) \circ L_k^1 \circ g_1(\Delta_1) [b_0 \otimes b_1] &= \sum_{\substack{r_0, r_1, \\ y, y_1}} f_1(r_0, y) \sqrt{r_1} g_1(y_1) b_0 E_{r_0} E_y^\Delta [E_{r_1} E_{y_1}^\Delta (b_1)] \\ &= \sum_{\substack{r_0, r_1, \\ y, y_1, \\ z, z_1}} f_1(r_0, y) \sqrt{r_1} g_1(y_1) b_0 E_{r_0} E_z E_{r_1} E_{z_1} b_1 E_{y_1 z_1} E_{yz} \end{aligned}$$

the projections products imply $r_0 = z = r_1 = z_1$ and $y_1 z_1 = yz$, so this is equal to

$$\begin{aligned} \sum_{\substack{r_0, y, \\ z}} f_1(r_0, y) \sqrt{r_0} g_1(y) b_0 E_{r_0} E_z b_1 E_{yz} &= \sum_{\substack{r_0, y, \\ z}} \sqrt{r_0} f_1(r_0, y) g_1(y) b_0 E_{r_0} E_y^\Delta (b_1) \\ &= \mathbf{m} \circ R_k^0 \circ f_1(R_u^0, \Delta_1) \circ g_1(\Delta_1) [b_0 \otimes b_1]. \end{aligned}$$

For the second relation, we compute the LHS on $b_0 \otimes b_1 \otimes b_2$:

$$\begin{aligned} \mathbf{m} \circ f_1(R_u^0, \Delta_1) \circ \mathbf{m}_{12} \circ L_k^1 \circ g_2(\Delta_1, \Delta_2)[b_0 \otimes b_1 \otimes b_2] \\ = \sum_{\substack{r_0, r_1, \\ y, y_1, y_2}} f_1(r_0, y) \sqrt{r_1} g_2(y_1, y_2) b_0 E_{r_0} E_y^\Delta [E_{r_1} E_{y_1}^\Delta (b_1) E_{y_2}^\Delta (b_2)] \\ = \sum_{\substack{r_0, r_1, \\ y, y_1, y_2, \\ z, z_1, z_2}} f_1(r_0, y) \sqrt{r_1} g_2(y_1, y_2) b_0 E_{r_0} E_z E_{r_1} E_{z_1} b_1 E_{y_1 z_1} E_{z_2} b_2 E_{y_2 z_2} E_{y z} \end{aligned}$$

which implies $r_0 = z = r_1 = z_1, y_1 z_1 = z_2$, and $y_2 z_2 = yz$, and

$$\begin{aligned} &= \sum_{\substack{r_0, z_1, z_2, \\ y_1, y_2}} f(r_0, y_1 y_2) \sqrt{r_0} g_2(y_1, y_2) b_0 E_{r_0} E_{z_1} b_1 E_{y_1 z_1} E_{z_2} b_2 E_{y_2 z_2} \\ &= \sum_{r_0, y_1, y_2} \sqrt{r_0} f(r_0, y_1 y_2) g_2(y_1, y_2) b_0 E_{r_0} E_{y_1}^\Delta (b_1) E_{y_2}^\Delta (b_2) \\ &= \mathbf{m} \circ R_k^0 \circ f(R_u^0, \Delta_1 \Delta_2) \circ g_2(\Delta_1, \Delta_2)[b_0 \otimes b_1 \otimes b_2]. \end{aligned}$$

For the third relation, we compute the LHS on $b_0 \otimes b_1 \otimes b_2$:

$$\begin{aligned} \mathbf{m} \circ f(R_u^0, \Delta_1, \Delta_2) \circ L_k^1 \circ L_k^2 \circ g_2(\Delta_1, \Delta_2)[b_0 \otimes b_1 \otimes b_2] \\ = \sum_{\substack{r_0, r_1, r_2, \\ y_1, y_2, y'_1, y'_2}} f_2(r_0, y_1, y_2) \sqrt{r_1} \sqrt{r_2} g_2(y'_1, y'_2) b_0 E_{r_0} E_{y_1}^\Delta [E_{r_1} E_{y'_1}^\Delta (b_1)] E_{y_2}^\Delta [E_{r_2} E_{y'_2}^\Delta (b_2)] \\ = \sum_{\substack{r_0, r_1, r_2, \\ y_1, y_2, y'_1, y'_2, \\ z_1, z_2, z'_1, z'_2}} f_2(r_0, y_1, y_2) \sqrt{r_1} \sqrt{r_2} g_2(y'_1, y'_2) b_0 E_{r_0} E_{z_1} E_{r_1} E_{z'_1} b_1 E_{y'_1 z'_1} E_{y_1 z_1} E_{z_2} E_{r_2} E_{z'_2} b_2 E_{y_2 z'_2} E_{y_2 z_2} \end{aligned}$$

which implies $r_0 = z_1 = r_1 = z'_1, y'_1 z'_1 = y_1 z_1 = z_2 = r_2 = z'_2$, and $y'_2 z'_2 = y_2 z_2$, so that:

$$\begin{aligned} &= \sum_{\substack{r_0, z_1, z_2, \\ y_1, y_2}} f_2(r_0, y_1, y_2) \sqrt{r_0} \sqrt{y_1 r_0} g_2(y_1, y_2) b_0 E_{r_0} E_{z_1} b_1 E_{y_1 z_1} E_{z_2} b_2 E_{y_2 z_2} \\ &= \sum_{r_0, y_1, y_2} r_0 \sqrt{y_1} f_2(r_0, y_1, y_2) g_2(y_1, y_2) b_0 E_{r_0} E_{y_1}^\Delta (b_1) E_{y_2}^\Delta (b_2) \\ &= \mathbf{m} \circ R_u^0 \circ \Delta_1^{1/2} \circ f_2(R_u^0, \Delta_1, \Delta_2) \circ g_2(\Delta_1, \Delta_2)[b_0 \otimes b_1 \otimes b_2]. \quad \blacksquare \end{aligned}$$

We can now change (5.8) in order to compare with [8, Theorem 5.2]. As in Lemma B.1, let

$$f_{\Delta k}(r_0, y_1) := F_{\Delta k}(r_0, r_0 y_1), \quad f_{\partial k \partial k}^{\mu\nu}(r_0, y_1, y_2) := F_{\partial k \partial k}^{\mu\nu}(r_0, r_0 y_1, r_0 y_1 y_2).$$

Using Lemmas B.2 and B.3, one gets

$$\begin{aligned} \mathbf{m} \circ F_{\Delta k}(R_u^0, R_u^1)[a \otimes \Delta k] &= \mathbf{m} \circ f_{\Delta k}(R_u^0, \Delta_1)[a \otimes \Delta k] \\ &= \mathbf{m} \circ f_{\Delta k}(R_u^0, \Delta_1) \circ L_k^1 \circ g_1(\Delta_1)[a \otimes \Delta h] \\ &\quad - g^{\mu\nu} \mathbf{m} \circ f_{\Delta k}(R_u^0, \Delta_1) \circ L_k^1 \circ \mathbf{m}_{12} \circ g_2(\Delta_1, \Delta_2)[a \otimes (\delta_\mu h) \otimes (\delta_\nu h)] \\ &= \mathbf{m} \circ R_k^0 \circ f_{\Delta k}(R_u^0, \Delta_1) \circ g_1(\Delta_1)[a \otimes \Delta h] \\ &\quad - g^{\mu\nu} \mathbf{m} \circ R_k^0 \circ f_{\Delta k}(R_u^0, \Delta_1) \circ g_2(\Delta_1, \Delta_2)[a \otimes (\delta_\mu h) \otimes (\delta_\nu h)] \\ &= 2\mathbf{m} \circ R_k^0 \circ f_{\Delta k}(R_u^0, \Delta_1) \circ g_1(\Delta_1)[a \otimes \Delta \ln k] \\ &\quad - 4g^{\mu\nu} \mathbf{m} \circ R_k^0 \circ f_{\Delta k}(R_u^0, \Delta_1 \Delta_2) \circ g_2(\Delta_1, \Delta_2)[a \otimes (\delta_\mu \ln k) \otimes (\delta_\nu \ln k)] \end{aligned}$$

and

$$\begin{aligned} \mathbf{m} \circ F_{\partial k \partial k}^{\mu\nu}(R_u^0, R_u^1, R_u^2)[a \otimes (\delta_\mu k) \otimes (\delta_\nu k)] &= \mathbf{m} \circ f_{\partial k \partial k}^{\mu\nu}(R_u^0, \Delta_1, \Delta_2)[a \otimes (\delta_\mu k) \otimes (\delta_\nu k)] \\ &= \mathbf{m} \circ f_{\partial k \partial k}^{\mu\nu}(R_u^0, \Delta_1, \Delta_2) \circ L_k^1 \circ L_k^2 \circ g_1(\Delta_1) \circ g_1(\Delta_2)[a \otimes (\delta_\mu h) \otimes (\delta_\nu h)] \\ &= \mathbf{m} \circ R_u^0 \circ \Delta_1^{1/2} \circ f_{\partial k \partial k}^{\mu\nu}(R_u^0, \Delta_1, \Delta_2) \circ g_1(\Delta_1) \circ g_1(\Delta_2)[a \otimes (\delta_\mu h) \otimes (\delta_\nu h)] \\ &= 4\mathbf{m} \circ R_u^0 \circ \Delta_1^{1/2} \circ f_{\partial k \partial k}^{\mu\nu}(R_u^0, \Delta_1, \Delta_2) \circ g_1(\Delta_1) \circ g_1(\Delta_2)[a \otimes (\delta_\mu \ln k) \otimes (\delta_\nu \ln k)]. \end{aligned}$$

So, the sum gives

$$\begin{aligned} & \mathbf{m} \circ F_{\Delta k}(R_u^0, R_u^1) [a \otimes \Delta k] + \mathbf{m} \circ F_{\partial k \partial k}^{\mu\nu}(R_u^0, R_u^1, R_u^2) [a \otimes (\delta_\mu k) \otimes (\delta_\nu k)] \\ & = \mathbf{m} \circ G_{(\Delta \ln k)}(R_u^0, \Delta_1) [a \otimes \Delta \ln k] + \mathbf{m} \circ G_{(\partial \ln k)(\partial \ln k)}^{\mu\nu}(R_u^0, \Delta_1, \Delta_2) [a \otimes (\delta_\mu \ln k) \otimes (\delta_\nu \ln k)] \end{aligned}$$

with

$$\begin{aligned} G_{(\Delta \ln k)}(R_u^0, \Delta_1) & := 2R_k^0 \circ f_{\Delta k}(R_u^0, \Delta_1) \circ g_1(\Delta_1), \\ G_{(\partial \ln k)(\partial \ln k)}^{\mu\nu}(R_u^0, \Delta_1, \Delta_2) & := 4R_u^0 \circ \Delta_1^{1/2} \circ f_{\partial k \partial k}^{\mu\nu}(R_u^0, \Delta_1, \Delta_2) \circ g_1(\Delta_1) \circ g_1(\Delta_2) \\ & \quad - 4g^{\mu\nu} R_k^0 \circ f_{\Delta k}(R_u^0, \Delta_1 \Delta_2) \circ g_2(\Delta_1, \Delta_2). \end{aligned}$$

The associated spectral functions are

$$\begin{aligned} G_{(\Delta \ln k)}(r_0, y_1) & = 2\sqrt{r_0} f_{\Delta k}(r_0, y_1) g_1(y_1), \\ G_{(\partial \ln k)(\partial \ln k)}^{\mu\nu}(r_0, y_1, y_2) & = 4r_0 \sqrt{y_1} f_{\partial k \partial k}^{\mu\nu}(r_0, y_1, y_2) g_1(y_1) g_1(y_2) - 4g^{\mu\nu} \sqrt{r_0} f_{\Delta k}(r_0, y_1 y_2) g_2(y_1, y_2). \end{aligned}$$

Another change of convention concerns the derivations of $C^\infty(\mathbb{T}_\Theta^2)$: in [8], $\widehat{\delta}_\mu := -i\delta_\mu$ is used. This implies that their expressions like $(\widehat{\delta}_\mu \ln k)(\widehat{\delta}_\nu \ln k)$ correspond to our $-(\delta_\mu \ln k) \otimes (\delta_\nu \ln k)$. Notice also their combination $\widehat{\delta}_1^2 \ln k + |\tau|^2 \widehat{\delta}_2^2 \ln k + 2\tau_1 \tau_2 \widehat{\delta}_1 \widehat{\delta}_2 \ln k = g^{\mu\nu} \widehat{\delta}_\mu \widehat{\delta}_\nu \ln k = -g^{\mu\nu} \delta_\mu \delta_\nu \ln k = \Delta \ln k$. Thus for a comparison of the two results, a $-$ sign has to be taken into account for the $G_{(\partial \ln k)(\partial \ln k)}^{\mu\nu}$ term. Finally, [8, Theorem 5.2] is written in terms of functions of $\ln \Delta$, thus it remains to make the final change of variables $y_1 = e^x$ in $G_{(\Delta \ln k)}$, and $y_1 = e^s$ and $y_2 = e^t$ in $G_{(\partial \ln k)(\partial \ln k)}^{\mu\nu}$.

In [8, Theorem 5.2], $\mathcal{R}_{FK} = -\frac{\pi}{\tau_2} \times$ [expression in R_1, R_2, W] while in (5.8), one has written $\mathcal{R}_2 = \frac{1}{4\pi} \times$ [expression in $F_{\Delta k}, F_{\partial k \partial k}^{\mu\nu}$]. The proof that $\mathcal{R}_2 = \frac{\tau_2}{(2\pi)^2} \mathcal{R}_{FK}$ is then equivalent to check that [expression in R_1, R_2, W] = $-$ [expression in $F_{\Delta k}, F_{\partial k \partial k}^{\mu\nu}$]. The previous technical results imply that this is equivalent to show that

$$\begin{aligned} R_1(x) & = -G_{(\Delta \ln k)}(r_0, e^x), \\ R_2(s, t) & = G_{(\partial \ln k)(\partial \ln k)}^{11}(r_0, e^s, e^t), \\ |\tau|^2 R_2(s, t) & = G_{(\partial \ln k)(\partial \ln k)}^{22}(r_0, e^s, e^t), \\ \tau_1 R_2(s, t) - i\tau_2 W(s, t) & = G_{(\partial \ln k)(\partial \ln k)}^{12}(r_0, e^s, e^t), \\ \tau_1 R_2(s, t) + i\tau_2 W(s, t) & = G_{(\partial \ln k)(\partial \ln k)}^{21}(r_0, e^s, e^t). \end{aligned}$$

All these relations can be checked directly. In particular, the relations on the RHS are independent of the variable r_0 .

In order to compare (5.9) for the noncommutative four torus with [11, Theorem 5.4], we can use the results in [14]. As before, we need the correspondence (A.5) between our trace φ and their trace $\varphi_0 \equiv \iota$. Here $g_i^{\mu\nu} = \delta^{\mu\nu}$ on the base tori $\mathbb{T}_{B,i}^2$, so that $|g_i|^{1/2} = 1$. Denote by \mathcal{R}_{FK} the curvature obtained in [11, Theorem 5.4], which is π^2 times [11, eq. (5.1)]. A comparison between eq. (1) and (3) in [14] and [11, eq. (5.1)] gives

$$\begin{aligned} \varphi_0(a\mathcal{R}_{FK}) & = \frac{1}{2}\varphi_0 \left(a \int_{\mathbb{S}^3} b_2(\xi) d\Omega \right) = \frac{\pi^2}{2}\varphi_0 \left(a[-\delta^{\mu\nu} k^{-2} (\widehat{\delta}_\mu \widehat{\delta}_\nu k^2) k^{-2} + \frac{3}{2} \delta^{\mu\nu} k^{-2} (\widehat{\delta}_\mu k^2) k^{-2} (\widehat{\delta}_\nu k^2) k^{-2}] \right) \\ & = \frac{\pi^2}{2(2\pi)^4} \varphi \left(a[\delta^{\mu\nu} k^{-2} (\delta_\mu \delta_\nu k^2) k^{-2} - \frac{3}{2} \delta^{\mu\nu} k^{-2} (\delta_\mu k^2) k^{-2} (\delta_\nu k^2) k^{-2}] \right) = \varphi(a\mathcal{R}_2), \end{aligned}$$

and the two results coincide.²

Appendix C. Supplementary data

Supplementary material related to this article can be found online at <http://dx.doi.org/10.1016/j.geomphys.2018.02.014>.

References

- [1] B. Iochum, T. Masson, Heat trace for Laplace type operators with non-scalar symbols, *J. Geom. Phys.* 116 (2017) 90–118.
- [2] I.G. Avramidi, Gauged gravity via spectral asymptotics of non-Laplace type operators, *J. High Energy Phys.* 2004 (07) (2004) 030.
- [3] I.G. Avramidi, T.P. Branson, Heat kernel asymptotics of operators with non-Laplace principal part, *Rev. Math. Phys.* 13 (07) (2001) 847–890.
- [4] Wolfram Research Inc., *Mathematica*, Version 11.1, 2017.
- [5] B. Iochum, T. Masson, Heat asymptotics for nonminimal Laplace type operators and application to noncommutative tori, *Mathematica Notebook* added as ancillary file on arXiv, 2017.
- [6] A. Connes, P. Tretkoff, The Gauss–Bonnet theorem for the noncommutative two torus, in: *Noncommutative Geometry, Arithmetic, and Related Topics*, Johns Hopkins Univ. Press, Baltimore, MD, 2011, pp. 141–158.
- [7] A. Connes, H. Moscovici, Modular curvature for noncommutative two-tori, *J. Amer. Math. Soc.* 27 (3) (2014) 639–684.

² The constant c computed in [14, eq. (3)] is not $c = 1/(2\pi^2)$ as claimed but $c = 1/2$ as shown by a direct comparison between [14, eq. (3)] and [11, eq. (5.1)] where a factor π^2 is not written. This explains the factor $\frac{1}{2}$ in our first equality.

- [8] F. Fathizadeh, M. Khalkhali, Scalar curvature for the noncommutative two torus, *J. Noncommut. Geom.* 7 (2013) 1145–1183.
- [9] F. Fathizadeh, M. Khalkhali, The Gauss-Bonnet theorem for noncommutative two tori with a general conformal structure, *J. Noncommut. Geom.* 6 (2012) 457–480.
- [10] L. Dabrowski, A. Sitarz, Curved noncommutative torus and Gauss–Bonnet, *J. Math. Phys.* 54 (2013) 013518.
- [11] F. Fathizadeh, M. Khalkhali, Scalar curvature for noncommutative four-tori, *J. Noncommut. Geom.* 9 (2015) 473–503.
- [12] S. Azzali, C. Lévy, C. Neira-Jiménez, S. Paycha, Traces of holomorphic families of operators on the noncommutative torus and on Hilbert Modules, in: *Geometrics Methods in Physics*, Birkhäuser, 2014, pp. 3–38.
- [13] A. Sitarz, Wodzicki residue and minimal operators on a noncommutative 4-dimensional torus, *J. Pseudo-Diff. Oper. Appl.* 5 (2014) 305–317.
- [14] F. Fathizadeh, On the scalar curvature for the noncommutative four torus, *J. Math. Phys.* 56 (6) (2015) 062303.
- [15] L. Dabrowski, A. Sitarz, An asymmetric noncommutative torus, *SIGMA* 11 (2015) 075, 11 pages.
- [16] Y. Liu, Modular curvature for toric noncommutative manifolds, 2015, arXiv:1510.04668v2 [math.OA].
- [17] S. Sadeghi, On Logarithmic Sobolev Inequality and a Scalar Curvature Formula for Noncommutative Tori, (Ph.D. thesis), Western University, Ontario, 2016.
- [18] A. Connes, F. Fathizadeh, The term a_4 in the heat kernel expansion of noncommutative tori, 2016, arXiv:1611.09815v1 [math.QA].
- [19] A. Connes, C^* -algèbres et géométrie différentielle, *C. R. Acad. Sci. Paris A* 290 (1980) 599–604.
- [20] M. Lesch, H. Moscovici, Modular curvature and Morita equivalence, *Geom. Funct. Anal.* 26 (2016) 818–873.
- [21] G.K. Pedersen, On the operator equation $HT + TH = 2K$, *Indiana Univ. Math. J.* 25 (11) (1976) 1029–1033.
- [22] P.B. Gilkey, *Invariance Theory, the Heat Equation and the Atiyah-Singer Index Theorem*, second ed., in: *Studies in Advanced Mathematics*, CRC Press, Inc, 1995.
- [23] P.B. Gilkey, *Asymptotic Formulae in Spectral Geometry*, CRC Press, 2003.
- [24] J.M.G. Fell, The structure of algebras of operator fields, *Acta Math.* 106 (3–4) (1961) 233–280.
- [25] B. Blackadar, *Operator Algebras: Theory of C^* -Algebras and Von Neumann Algebras*, Vol. 122, Springer Science, 2006.
- [26] J.M. Gracia-Bondía, J.C. Várilly, H. Figueroa, *Elements of Noncommutative Geometry*, Birkhäuser, Boston, 2001.
- [27] M. Dubois-Violette, A. Kriegl, Y. Maeda, P.W. Michor, Smooth $*$ -algebras, *Progr. Theoret. Phys. Suppl.* 144 (2002) 54–78.
- [28] M. Lesch, Divided differences in noncommutative geometry: Rearrangement lemma, functional calculus and expansional formula, *J. Noncommut. Geom.* 11 (2017) 193–223.